



National Comprehensive  
Cancer Network®

NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®)

# Small Bowel Adenocarcinoma

Version 2.2021 — September 10, 2021

[NCCN.org](https://www.nccn.org)

[Continue](#)



**\*Al B. Benson, III, MD/Chair †**  
Robert H. Lurie Comprehensive Cancer  
Center of Northwestern University

**\*Alan P. Venook, MD/Vice-Chair † ‡**  
UCSF Helen Diller Family  
Comprehensive Cancer Center

**Mahmoud M. Al-Hawary, MD †**  
University of Michigan Rogel Cancer Center

**Nilofer Azad, MD †**  
The Sidney Kimmel Comprehensive  
Cancer Center at Johns Hopkins

**\*Yi-Jen Chen, MD, PhD §**  
City of Hope National Medical Center

**Kristen K. Ciombor, MD †**  
Vanderbilt-Ingram Cancer Center

**Stacey Cohen, MD †**  
Fred Hutchinson Cancer Research Center/  
Seattle Cancer Care Alliance

**Harry S. Cooper, MD ≠**  
Fox Chase Cancer Center

**Dustin Deming, MD †**  
University of Wisconsin Carbone Cancer Center

**Linda Farkas, MD ¶**  
UT Southwestern Simmons  
Comprehensive Cancer Center

**Ignacio Garrido-Laguna, MD, PhD †**  
Huntsman Cancer Institute  
at the University of Utah

**Jean L. Grem, MD †**  
Fred & Pamela Buffett Cancer Center

**Andrew Gunn, MD †**  
O'Neal Comprehensive Cancer Center at UAB

**J. Randolph Hecht, MD †**  
UCLA Jonsson Comprehensive Cancer Center

**Sarah Hoffe, MD §**  
Moffitt Cancer Center

**Joleen Hubbard, MD ‡**  
Mayo Clinic Cancer Center

**Steven Hunt, MD ¶**  
Siteman Cancer Center at Barnes-  
Jewish Hospital and Washington  
University School of Medicine

**Kimberly L. Johung, MD, PhD §**  
Yale Cancer Center/Smilow Cancer Hospital

**Natalie Kirilcuk, MD ¶**  
Stanford Cancer Institute

**Smitha Krishnamurthi, MD † †**  
Case Comprehensive Cancer Center/  
University Hospitals Seidman Cancer Center  
and Cleveland Clinic Taussig Cancer Institute

**Wells A. Messersmith, MD †**  
University of Colorado Cancer Center

**\*Jeffrey Meyerhardt, MD, MPH †**  
Dana-Farber Brigham and Women's  
Cancer Center

**\*Eric D. Miller, MD, PhD §**  
The Ohio State University Comprehensive  
Cancer Center - James Cancer Hospital  
and Solove Research Institute

**Mary F. Mulcahy, MD ‡ †**  
Robert H. Lurie Comprehensive Cancer  
Center of Northwestern University

**Steven Nurkin, MD, MS ¶**  
Roswell Park Comprehensive Cancer Center

**Michael J. Overman, MD † ‡**  
The University of Texas  
MD Anderson Cancer Center

**Aparna Parikh, MD †**  
Massachusetts General Hospital Cancer Center

**Hitendra Patel, MD †**  
UC San Diego Moores Cancer Center

**Katrina Pedersen, MD, MS †**  
Siteman Cancer Center at Barnes-  
Jewish Hospital and Washington  
University School of Medicine

**Elizabeth Raskin, MD ¶**  
UC Davis Comprehensive Cancer Center

**\*Leonard Saltz, MD † ‡ †**  
Memorial Sloan Kettering Cancer Center

**Charles Schneider, MD †**  
Abramson Cancer Center  
at the University of Pennsylvania

**David Shibata, MD ¶**  
The University of Tennessee  
Health Science Center

**John M. Skibber, MD ¶**  
The University of Texas  
MD Anderson Cancer Center

**\*Constantinos T. Sofocleous, MD, PhD †**  
Memorial Sloan Kettering Cancer Center

**Elena M. Stoffel, MD, MPH ≠**  
University of Michigan Rogel Cancer Center

**Eden Stotsky-Himelfarb, BSN, RN † ¶ ≠**  
The Sidney Kimmel Comprehensive  
Cancer Center at Johns Hopkins

**\*Christopher G. Willett, MD §**  
Duke Cancer Institute

**NCCN**  
**Kristina Gregory, RN, MSN**  
**Lisa Gurski, PhD**

**Continue**

† Diagnostic/Interventional radiology	≠ Pathology
≠ Gastroenterology	¶ Patient advocate
‡ Hematology/Hematology oncology	§ Radiotherapy/Radiation oncology
¶ Surgery/Surgical oncology	¶ Surgery/Surgical oncology
‡ Internal medicine	* Discussion Section Writing Committee
† Medical oncology	



### SUBCOMMITTEE

**Katrina Pedersen, MD, MS †/Lead**

Siteman Cancer Center at Barnes-Jewish Hospital  
and Washington University School of Medicine

**Mustafa A. Arain, MD ☒**

UCSF Helen Diller Family Comprehensive Cancer Center

**Al B. Benson, III, MD/Chair †**

Robert H. Lurie Comprehensive Cancer Center of Northwestern University

**Michael J. Overman, MD † ‡**

The University of Texas MD Anderson Cancer Center

**John M. Skibber, MD ¶**

The University of Texas MD Anderson Cancer Center

**Alan P. Venook, MD/Vice-Chair † ‡**

UCSF Helen Diller Family Comprehensive Cancer Center

[NCCN Guidelines Panel Disclosures](#)

**Continue**

☒ Gastroenterology  
‡ Hematology/Hematology oncology  
† Medical oncology  
¶ Surgery/Surgical oncology



### [NCCN Small Bowel Adenocarcinoma Panel Members](#) [Summary of the Guidelines Updates](#)

#### [Duodenum](#)

- [Workup and Primary Treatment \(SBA-1\)](#)
- [Adjuvant Therapy \(SBA-2\)](#)

#### [Jejunum/Ileum](#)

- [Workup and Primary Treatment \(SBA-3\)](#)
- [Adjuvant Therapy \(SBA-4\)](#)

#### [Metastatic Adenocarcinoma \(SBA-5\)](#) [Recurrence \(SBA-5\)](#)

#### [Principles of Imaging and Endoscopy \(SBA-A\)](#)

#### [Principles of Pathologic Review \(SBA-B\)](#)

#### [Principles of Surgery \(SBA-C\)](#)

#### [Principles of Systemic Therapy \(SBA-D\)](#)

#### [Principles of Radiation Therapy \(SBA-E\)](#)

#### [Principles of Survivorship \(SBA-F\)](#)

#### [Staging \(ST-1\)](#)

**Clinical Trials:** NCCN believes that the best management for any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

To find clinical trials online at NCCN Member Institutions, [click here: nccn.org/clinical\\_trials/member\\_institutions.aspx](#).

**NCCN Categories of Evidence and Consensus:** All recommendations are category 2A unless otherwise indicated.

See [NCCN Categories of Evidence and Consensus](#).

The NCCN Guidelines® are a statement of evidence and consensus of the authors regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult the NCCN Guidelines is expected to use independent medical judgment in the context of individual clinical circumstances to determine any patient's care or treatment. The National Comprehensive Cancer Network® (NCCN®) makes no representations or warranties of any kind regarding their content, use or application and disclaims any responsibility for their application or use in any way. The NCCN Guidelines are copyrighted by National Comprehensive Cancer Network®. All rights reserved. The NCCN Guidelines and the illustrations herein may not be reproduced in any form without the express written permission of NCCN. ©2021.



Updates in Version 2.2021 of the NCCN Guidelines for Small Bowel Adenocarcinoma from Version 1.2021 include:

#### **SBA-D Systemic Therapy for Advanced or Metastatic Disease**

- Dostarlimab-gxly added as a subsequent therapy option for dMMR/MSI-H (SBA-D 1 and SBA-D 2)
- Dostarlimab-gxly dosing (SBA-D 5) and reference (SBA-D 7) added.

Updates in Version 1.2021 of the NCCN Guidelines for Small Bowel Adenocarcinoma from Version 2.2020 include:

#### **SBA-1**

- Resectable disease
  - ▶ Wording changed to Resection with en bloc removal of regional lymph nodes (also applied to SBA-3)
  - ▶ The specific surgical procedures removed, as these are noted on SBA-C. Resection now listed. (also applies to SBA-3)
- The following added:
  - ▶ Advanced ampullary cancer
    - ◊ Consider systemic therapy (SBA-D) as per the NCCN Guidelines for Small Bowel Adenocarcinoma

#### **SBA-2**

- Adjuvant Treatment
  - ▶ T3, N0, M0 with high-risk features or T4, N0, M0 (MSS or pMMR) or T Any, N1–2
    - ◊ Duration of therapy for FOLFOX or CAPEOX changed from 6 mo to 3-6 mo (also applies to SBA-4)
  - ▶ Footnote k modified: ~~The IDEA trial, which successfully showed non-inferior 3-year disease-free survival with 3 months of CAPEOX compared to 6 months of CAPEOX enrolled no patients with SBAs, which tend to have a higher risk for recurrence when compared to colon cancer. As a result, data extrapolation is not recommended for SBA patients receiving adjuvant therapy. No patients with SBA were included in the IDEA pooled analysis of adjuvant colon cancer trials. However, in the absence of any direct data regarding SBA, the finding of non-inferior 3-year disease-free survival with 3 months of CAPEOX compared to 6 months of CAPEOX in colon cancer may be extrapolated.~~

#### **SBA-A 1 of 2**

- Initial Workup/Staging; Chest, abdomen, pelvic CT
  - ▶ Sub-bullet 6 modified: ~~If IV iodinated contrast is contraindicated, then MR examination of the abdomen/pelvis with IV gadolinium-based contrast may be obtained instead. Gadolinium-based contrast is contraindicated in patients on dialysis.~~ In patients with chronic kidney failure (*including patients on dialysis*) ~~stage IV who are not on dialysis~~, IV gadolinium-based contrast may be administered in select cases using ~~gadofosveset trisodium, gadoxetate disodium, gadobutrol, gadopentetate dimeglumine, gadobenate dimeglumine, or gadoteridol.~~
  - ▶ Reference 2 added.

#### **SBA-B**

- Pathologic stage
  - ▶ Bullet 1; sub-bullet 8 modified: ~~MSI/DNA-MMR deficiency~~ status
- Microsatellite Instability/Mismatch Repair Testing
  - ▶ Bullet 1 modified: Universal MMR or MSI testing is recommended in all *newly diagnosed* patients with ~~a personal history of~~ small bowel adenocarcinoma

#### **SBA-D 1 of 7**

- Initial Therapy
  - ▶ The following treatment option added
    - ◊ (Nivolumab ± ipilimumab) or pembrolizumab (dMMR/MSI-H only)
  - ▶ Footnote b added: For infection risk, monitoring, and prophylaxis recommendations for targeted therapies, see INF-A in the NCCN Guidelines for Prevention and Treatment of Cancer-Related Infections. (also applies to SBA-D 2 of 7)
  - ▶ Footnote d modified: Bevacizumab had been shown to be safe in advanced SBA, though efficacy has not been proven. *An FDA-approved biosimilar is an appropriate substitute for bevacizumab.*
  - ▶ Footnote e added: See NCCN Guidelines for Management of Immunotherapy-Related Toxicities. (also applies to SBA-D 2 of 7)
  - ▶ Footnote f added: If no previous treatment with a checkpoint inhibitor. (also applies to SBA-D 2 of 7)
- Subsequent Therapy
  - ▶ Footnote g modified: Larotrectinib or *entrectinib* are treatment options for patients with metastatic SBA that is *NTRK* gene fusion positive. (also applies to SBA-D 2 of 7)

#### **SBA-D 4 of 7**

- FOLFOXIRI dosing modified with dosage reduction for fluorouracil: Fluorouracil ~~4600~~1200 mg/m<sup>2</sup>/day x 2 days (total ~~3200~~2400 mg/m<sup>2</sup> over 48 hours)


#### **SBA-D 6 of 7**

- Albumin-bound paclitaxel dosing modified to provide a range 220–260 mg/m<sup>2</sup>
- The following dosing regimen added with reference
  - ▶ Entrectinib (*NTRK* gene fusion positive)
    - ◊ 600 mg PO once daily



# NCCN Guidelines Version 2.2021

## Small Bowel Adenocarcinoma



<sup>a</sup> All patients with small bowel adenocarcinoma (SBA) should be counseled for familial malignancies and considered for risk assessment, including Lynch syndrome (hereditary nonpolyposis colorectal cancer [HNPCC]), familial adenomatous polyposis (FAP), and other polyloid mutations. Refer to the [NCCN Guidelines for Genetic/Familial High-Risk Assessment: Colorectal](#).  
<sup>b</sup> See [Principles of Imaging and Endoscopy \(SBA-A\)](#).  
<sup>c</sup> EUS should be considered when needed to discern duodenal malignancy from ampullary, distal common bile duct, or pancreatic head malignancy. Also consider if other radiologic imaging is insufficient for clinical staging.

<sup>d</sup> See [Principles of Pathologic Review \(SBA-B\)](#). Depending on tumor location and patient history, celiac disease or Crohn's disease may need to be assessed.  
<sup>e</sup> See [Principles of Surgery \(SBA-C\)](#).  
<sup>f</sup> See [Principles of Radiation Therapy \(SBA-E\)](#).  
<sup>g</sup> Preoperative chemoradiation should be considered in patients who remain unresectable following a course of induction chemotherapy.

**Note: All recommendations are category 2A unless otherwise indicated.**  
**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**



# NCCN Guidelines Version 2.2021

## Small Bowel Adenocarcinoma

LOCATION	PATHOLOGIC STAGE <sup>d</sup>	ADJUVANT TREATMENT <sup>i,j</sup>	SURVEILLANCE <sup>m</sup>
Duodenum	T1–2, N0, M0; T3–4 N0, M0 (MSI-H or dMMR)	Observation	<ul style="list-style-type: none"> <li>• History and physical examination every 3–6 months for 2 years, then every 6 months for a total of 5 years</li> <li>• CEA and/or CA 19-9 every 3–6 months for 2 years then every 6 months for a total of 5 years</li> <li>• Chest/abdominal/pelvic CT every 6–12 months for 2 years, then every 12 months years 3–5</li> <li>• PET/CT is not indicated</li> <li>• Routine capsule endoscopy is not indicated</li> </ul> <p><a href="#">See Principles of Survivorship (SBA-F)</a> <a href="#">See Recurrence (SBA-5)</a></p>
	T3, N0, M0 (MSS or pMMR and no high-risk features)	Observation or Consider 5-FU/LV or capecitabine	
	T3, N0, M0 with high-risk features <sup>h</sup> or T4, N0, M0 (MSS or pMMR)	Observation or FOLFOX or CAPEOX (3–6 mo) <sup>k</sup> or 5-FU/LV or capecitabine (6 mo) <sup>l</sup> or Chemoradiation <sup>f</sup> with capecitabine or infusional 5-FU (only if positive margin)	
	T Any, N1–2	FOLFOX or CAPEOX (3–6 mo) <sup>k</sup> or 5-FU/LV or capecitabine (6 mo) <sup>l</sup> or Chemoradiation <sup>f</sup> with capecitabine or infusional 5-FU (only if positive margin)	

<sup>d</sup> See Principles of Pathologic Review (SBA-B). Depending on tumor location and patient history, celiac disease or Crohn's disease may need to be assessed.

<sup>f</sup> See Principles of Radiation Therapy (SBA-E).

<sup>h</sup> High-risk features in stage II SBA include close or positive resection margins, <5 lymph nodes examined if duodenal location or <8 lymph nodes examined if jejunal/ileal primary tumor location, and tumor perforation. Further consideration may be made for administering chemotherapy in patients with stage II disease who have lymphovascular or perineural invasion, or poorly differentiated histology due to data extrapolated from colorectal cancer studies.

<sup>i</sup> Enrollment in a clinical trial is encouraged [eg, Phase III Trial Investigating the Potential Benefit of Adjuvant Chemotherapy for Small Bowel Adenocarcinoma (BALLAD): <https://clinicaltrials.gov/ct2/show/NCT02502370>].

<sup>j</sup> See Principles of Systemic Therapy (SBA-D 3 of 7).

<sup>k</sup> No patients with SBA were included in the IDEA pooled analysis of adjuvant colon cancer trials. However, in the absence of any direct data regarding SBA, the finding of non-inferior 3-year disease-free survival with 3 months of CAPEOX compared to 6 months of CAPEOX in colon cancer may be extrapolated.

<sup>l</sup> Survival benefit in adding oxaliplatin to fluoropyrimidine has not been demonstrated in geriatric patients (>70 years) for colon cancer adjuvant management.

<sup>m</sup> No studies have been performed to assess ideal surveillance intervals for SBA. The data in colorectal cancer surveillance is generally accepted as appropriate for SBA.


**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 2.2021

## Small Bowel Adenocarcinoma



<sup>a</sup> All patients with SBA should be counseled for familial malignancies and considered for risk assessment, including Lynch syndrome (HNPCC), FAP, and other inherited polypoid mutations. Refer to the [NCCN Guidelines for Genetic/Familial High-Risk Assessment: Colorectal](#).

<sup>b</sup> [See Principles of Imaging and Endoscopy \(SBA-A\)](#).

<sup>d</sup> [See Principles of Pathologic Review \(SBA-B\)](#). Depending on tumor location and patient history, celiac disease or Crohn's disease may need to be assessed.

<sup>e</sup> [See Principles of Surgery \(SBA-C\)](#).

**Note:** All recommendations are category 2A unless otherwise indicated.


**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.





# NCCN Guidelines Version 2.2021

## Small Bowel Adenocarcinoma



<sup>d</sup> See [Principles of Pathologic Review \(SBA-B\)](#). Depending on tumor location and patient history, celiac disease or Crohn's disease may need to be assessed.

<sup>h</sup> High-risk features in stage II SBA include close or positive resection margins, <5 lymph nodes examined if duodenal location or <8 lymph nodes examined if jejunal/ileal primary tumor location, and tumor perforation. Further consideration may be made for administering chemotherapy in patients with stage II disease who have lymphovascular or perineural invasion, or poorly differentiated histology due to data extrapolated from colorectal cancer studies.

<sup>i</sup> Enrollment in a clinical trial is encouraged [eg, Phase III Trial Investigating the Potential Benefit of Adjuvant Chemotherapy for Small Bowel Adenocarcinoma (BALLAD): <https://clinicaltrials.gov/ct2/show/NCT02502370>].

<sup>j</sup> See [Principles of Systemic Therapy \(SBA-D 3 of 7\)](#).

<sup>k</sup> No patients with SBA were included in the IDEA pooled analysis of adjuvant colon cancer trials. However, in the absence of any direct data regarding SBA, the finding of non-inferior 3-year disease-free survival with 3 months of CAPEOX compared to 6 months of CAPEOX in colon cancer may be extrapolated.

<sup>l</sup> Survival benefit in adding oxaliplatin to fluoropyrimidine has not been demonstrated in geriatric patients (>70 years) for colon cancer adjuvant management.


<sup>m</sup> No studies have been performed to assess ideal surveillance intervals for SBA. The data in colorectal cancer surveillance is generally accepted as appropriate for SBA.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

### METASTATIC AND/OR RECURRENCE

### WORKUP



<sup>b</sup> See Principles of Imaging and Endoscopy ([SBA-A](#)).

<sup>d</sup> See Principles of Pathologic Review ([SBA-B](#)). Depending on location and patient history, celiac disease or Crohn’s disease may need to be assessed.

<sup>n</sup> Potentially resectable visceral or peritoneal metastases are extremely rare for SBA. See [Discussion](#) for information on metastastectomy and cytoreductive surgery/ intraperitoneal chemotherapy, which may be considered for select patients.

**Note: All recommendations are category 2A unless otherwise indicated.**  
**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**

**PRINCIPLES OF IMAGING AND ENDOSCOPY****Initial Workup/Staging****• Chest, abdomen, pelvic CT**

- ▶ Evaluate local extent of tumor infiltration into surrounding structures.
- ▶ Assess for distant metastatic disease to lungs, lymph node, liver, peritoneal cavity, and other organs.
- ▶ CT abdomen/pelvis should be performed with intravenous iodinated contrast and oral contrast agents unless contraindicated.
- ▶ CT chest does not require contrast (though may be given if performed with abdominal CT).
- ▶ Consider CT enterography or enteroclysis for specific cases where primary tumor is poorly visualized by standard methods.<sup>1</sup>
- ▶ If IV iodinated contrast is contraindicated, then MR examination of the abdomen/pelvis with IV gadolinium-based contrast may be obtained instead. In patients with chronic kidney failure (including patients on dialysis), IV gadolinium-based contrast may be administered in select cases using gadobutrol, gadopentetate dimeglumine, gadobenate dimeglumine, or gadoteridol.<sup>2</sup>

**• MRI**

- ▶ MR of the abdomen/pelvis may be considered where there is contraindication to CT.<sup>3</sup> MR enterography or enteroclysis may similarly be considered when conventional CT or MR with contrast have failed to discern a tumor.<sup>4,5</sup>
- ▶ Consider MR of the abdomen with contrast for further evaluation of indeterminate liver lesions on CT scan.
- ▶ Magnetic resonance cholangiopancreatography (MRCP) may need to be obtained in the initial workup of suspected duodenal malignancies to further ascertain tumor site of origin, particularly in cases of biliary obstruction.

**• PET/CT is not indicated as efficacy and clinical benefit have not been examined compared to CT or MR.<sup>6</sup>**

- ▶ Consider obtaining PET/CT in instances of equivocal CT or MR results, including the evaluation of potential peritoneal disease, particularly where potential lesions are sized greater than the lower limits of PET detection.

<sup>1</sup> Boudiaf M, Jaff A, Soyer P, et al. Small-bowel diseases: prospective evaluation of multi-detector row helical CT enteroclysis in 107 consecutive patients. *Radiology* 2004;233:338-344.

<sup>2</sup> Weinreb JC, Rodby RA, Yee J, et al. Use of intravenous gadolinium-based contrast media in patients with kidney disease: consensus statements from the American College of Radiology and the National Kidney Foundation. *Radiology* 2021;298:28-35.

<sup>3</sup> Masselli G, Casciani E, Poletti E, et al. Magnetic resonance imaging of small bowel neoplasms. *Cancer Imaging* 2013;13:92-99.

<sup>4</sup> Cronin CG, Lohan DG, Browne AM, et al. Magnetic resonance enterography in the evaluation of the small bowel. *Semin Roentgenol* 2009;44:237-243.

<sup>5</sup> Masselli G, Di Tola M, Casciani E, et al. Diagnosis of small-bowel diseases: Prospective comparison of multi-detector row CT enterography with MR enterography. *Radiology* 2016;279:420-431.

<sup>6</sup> Cronin CG, Scott J, Kambadakone A, et al. Utility of positron emission tomography/CT in the evaluation of small bowel pathology. *Br J Radiol* 2012;85:1211-1221.

**Note: All recommendations are category 2A unless otherwise indicated.**

**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**

**PRINCIPLES OF IMAGING AND ENDOSCOPY****Initial Workup/Staging****• Endoscopy****▶ EGD**

- ◊ May be used for detection and pathologic sampling where duodenal malignancy is suspected. If simultaneous intestinal obstruction is detected, palliative stenting may be considered.<sup>7</sup>

**▶ EUS**

- ◊ Useful for enhanced pre-therapeutic clinical staging of proximal small intestinal malignancies and to discern duodenal from ampullary, biliary, or pancreatic primaries.<sup>8</sup>

**▶ Push- or device-assisted enteroscopy (double- or single-balloon enteroscopy)**

- ◊ Not required for routine staging workup, though may be considered in patients with small intestinal strictures for diagnostic and/or palliative benefit. Biopsy may be performed during these procedures.<sup>9–11</sup>

**▶ Capsule endoscopy**

- ◊ Consider when radiographic imaging and other forms of endoscopy fail to reveal a suspected primary lesion.<sup>12,13</sup> This is not the preferred primary method for diagnostic workup due to inability to obtain tissue for diagnosis.
- ◊ Contraindicated where small bowel obstruction or strictures exist.

**Monitoring****• Chest, abdomen, pelvis CT with contrast**

- ▶ Prior to adjuvant therapy to assess response to resection or primary therapy.
- ▶ During re-evaluation of conversion to resectable disease (neoadjuvant therapy).
- ▶ CT enterography is not routinely indicated for monitoring and should be reserved for instances of clinical necessity.

**Surveillance**

- Surveillance for recurrence should be judiciously applied. Recommend similar approach to colorectal surveillance due to lack of small bowel adenocarcinoma (SBA)-specific data.
- Patients with confirmed Lynch syndrome should have appropriate screenings commensurate with their genotype and family cancer history. This may include small intestinal screening moving forward.<sup>14</sup>

<sup>7</sup> Hara AK, Leighton JA, Sharma VK, et al. Imaging of small bowel disease: comparison of capsule endoscopy, standard endoscopy, barium examination, and CT. *Radiographics* 2005;25:697-711; discussion 711-8.

<sup>8</sup> Nylund K, Odegaard S, Hausken T, et al. Sonography of the small intestine. *World J Gastroenterol* 2009;15:1319-1330.

<sup>9</sup> Cazzato IA, Cammarota G, Nista EC, et al. Diagnostic and therapeutic impact of double-balloon enteroscopy (DBE) in a series of 100 patients with suspected small bowel diseases. *Dig Liver Dis* 2007;39:483-487.

<sup>10</sup> Sunada K, Yamamoto H, Kita H, et al. Clinical outcomes of enteroscopy using the double-balloon method for strictures of the small intestine. *World J Gastroenterol* 2005;11:1087-1089.

<sup>11</sup> Chen WG, Shan GD, Zhang H, et al. Double-balloon enteroscopy in small bowel diseases: Eight years single-center experience in China. *Medicine (Baltimore)* 2016;95:e5104.

<sup>12</sup> Bailey AA, Debinski HS, Appleyard MN, et al. Diagnosis and outcome of small bowel tumors found by capsule endoscopy: a three-center Australian experience. *Am J Gastroenterol* 2006;101:2237-2243.

<sup>13</sup> Cobrin GM, Pittman RH, Lewis BS. Increased diagnostic yield of small bowel tumors with capsule endoscopy. *Cancer* 2006;107:22-27.

<sup>14</sup> Koornstra JJ. Small bowel endoscopy in familial adenomatous polyposis and Lynch syndrome. *Best Pract Res Clin Gastroenterol* 2012;26:359-368.

**Note: All recommendations are category 2A unless otherwise indicated.**

**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**

**PRINCIPLES OF PATHOLOGIC REVIEW****Small Bowel Adenocarcinoma Appropriate for Resection**

- Prefer pathologic confirmation of disease prior to resection, where possible

**Discrimination of Duodenal Malignancy from Other Primary Sites**

- To differentiate SBA from ampullary malignancies in duodenal cancers of the second portion, the epicenter of the tumor or precursor lesion should not be at the ampulla, and >75% of the mass should not be within the ampulla

**Pathologic Stage**

- The following parameters should be reported:
  - ▶ Primary tumor site (duodenum, jejunum, ileum, overlapping, not otherwise specified [NOS])
  - ▶ Grade of cancer
  - ▶ Tumor depth of invasion (T stage)
  - ▶ Number of lymph nodes evaluated and number of lymph nodes positive (N)
  - ▶ Status of proximal, distal, radial or mesenteric, uncinata, bile duct, pancreas, and other margins, as appropriate
  - ▶ Specimen orientation and inking involves both the pathologist and surgeon as this will help to ensure accurate assessment of the size and extent of the tumor. There should be either direct communication between the surgeon and pathologist for proper orientation and margin identification, or the surgeon should identify the important margins with a clearly understood and documented method (eg, written on the pathology requisition); see above – Status of margins.
  - ▶ Lymphovascular invasion
  - ▶ MSI/MMR status
  - ▶ Evidence of celiac disease
  - ▶ Presence of Crohn's disease
  - ▶ Presence of polyps

**Lymph Node Evaluation**

- The AJCC recommends a minimum evaluation of 8 lymph nodes, though the College of American Pathologists notes no clear number of minimum lymph nodes to predict complete lymph node negativity has been established
- Regional lymph nodes differ by site of primary tumor:
  - ▶ Duodenum: retropancreatic, hepatic artery, inferior pancreaticoduodenal, and superior mesenteric
  - ▶ Jejunum/ileum: cecal (terminal ileum only), ileocolic (terminal ileum only), superior mesenteric, mesenteric, and NOS

**Microsatellite Instability/Mismatch Repair Testing**

- Universal MMR or MSI testing is recommended in all newly diagnosed patients with SBA
- Incidence of deficient MMR (dMMR)/MSI-high (MSI-H) and the possibility of germline mutation is enriched in SBA patients compared to those with colon and rectal cancer, making this an important prognostic and/or predictive biomarker<sup>1,2</sup>
- Stage II dMMR/MSI-H patients may have improved survival compared to proficient MMR (pMMR)/microsatellite stable (MSS) patients; however, this has not been confirmed in the SBA population and is extrapolated from colorectal cancer data
- MMR or MSI testing should be performed only in CLIA-certified laboratories
- Testing for MSI may be performed by validated next-generation sequencing (NGS) panels

<sup>1</sup> Aparicio T, Svrcek M, Zaanani A, et al. Small bowel adenocarcinoma phenotyping, a clinicobiological prognostic study. *Br J Cancer* 2013;109:3057-3066.

<sup>2</sup> Schrock AB, Devoe CE, McWilliams R, et al. Genomic profiling of small-bowel adenocarcinoma. *JAMA Oncol* 2017;3:1546-1553.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PRINCIPLES OF SURGERY****Principles for All Primary Sites:**

- Intraoperative staging of the abdomen, particularly including the mesentery, omentum, and peritoneum, should be completed in all cases.
- Adequate lymph node dissection, consisting of the resection and evaluation of at least 8 lymph nodes, should be the goal for all resections.<sup>1,2</sup>

**Duodenum**

- **Pancreaticoduodenectomy (Whipple)**
  - ▶ Should be considered for all duodenal malignancies, particularly if arising in the second portion of the duodenum or invading any portion of the ampulla or pancreas.
  - ▶ Pylorus preservation is acceptable in the absence of a hereditary condition.
  - ▶ Minimally invasive procedures should be employed only by experienced surgeons.
  - ▶ Margins should be considered for frozen section at the time of resection if there are concerns about the margins. If margins <5 mm, the surgeon should consider re-excision of involved margin.
- **Limited segmentectomy**
  - ▶ Though controversial, may be considered for select cases in the absence of a hereditary condition, particularly in lesions on the anti-mesenteric side of the intestine that involve the third and fourth segments of the duodenum.
  - ▶ Due to reported lower yield of lymph nodes sampled during segmental resection, particular attention will need to be made for complete lymph node dissection and evaluation.<sup>3</sup>
  - ▶ Case reports suggest segmentectomy and other limited resection methods may be considered for lesions of the first portion of the duodenum,<sup>4</sup> particularly for lesions located on the mesenteric side of the intestine and for those <2 cm in size.

<sup>1</sup> Overman MJ, Hu CY, Kopetz S, et al. A population-based comparison of adenocarcinoma of the large and small intestine: insights into a rare disease. *Ann Surg Oncol* 2012;19:1439-1445.

**Jejunum/Ileum**

- **Segmentectomy**
  - ▶ Lymph nodes should be identified and resected down to the origin of feeder vessels. Clinically suspicious nodes outside of the field of resection should be biopsied or resected whenever possible.
  - ▶ Margins of at least 5–10 cm on either side of the tumor should be obtained.
  - ▶ Terminal ileal resection with right hemicolectomy should be used for distal ileal tumors.

<sup>2</sup> Overman MJ, Hu CY, Wolff RA, Chang GJ. Prognostic value of lymph node evaluation in small bowel adenocarcinoma: analysis of the surveillance, epidemiology, and end results database. *Cancer* 2010;116:5374-5382.

<sup>3</sup> Onkendi EO, Boostrom SY, Sarr MG, et al. 15-year experience with surgical treatment of duodenal carcinoma: a comparison of periampullary and extra-ampullary duodenal carcinomas. *J Gastrointest Surg* 2012;16:682-691.

<sup>4</sup> Hashimoto D, Arima K, Chikamoto A, et al. Limited resection of the duodenum for nonampullary duodenal tumors, with review of the literature. *Am Surg* 2016;82:1126-1132.


**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



# NCCN Guidelines Version 2.2021

## Small Bowel Adenocarcinoma



**Patient with prior oxaliplatin exposure or contraindication ([SBA-D 2 of 7](#))**  
**Regimen Dosing ([SBA-D 4 of 7](#))**

<sup>a</sup> Many of the regimens recommended in these guidelines are extrapolated from data for colorectal cancer.  
<sup>b</sup> For infection risk, monitoring, and prophylaxis recommendations for targeted therapies, see INF-A in the [NCCN Guidelines for Prevention and Treatment of Cancer-Related Infections](#).  
<sup>c</sup> For elderly patients, please complete geriatric assessment to aid appropriate prediction of treatment risks. [See NCCN Guidelines for Older Adult Oncology, OAO-2](#).  
<sup>d</sup> Bevacizumab has been shown to be safe in advanced SBA, though efficacy has not been proven. An FDA-approved biosimilar is an appropriate substitute for bevacizumab.  
<sup>e</sup> [See NCCN Guidelines for Management of Immunotherapy-Related Toxicities](#).  
<sup>f</sup> If no previous treatment with a checkpoint inhibitor.  
<sup>g</sup> Larotrectinib or entrectinib are treatment options for patients with metastatic SBA that is *NTRK* gene fusion positive.


**Note: All recommendations are category 2A unless otherwise indicated.**  
**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**

### PRINCIPLES OF SYSTEMIC THERAPY FOR ADVANCED OR METASTATIC DISEASE<sup>a,b</sup>

**PATIENT STATUS**

**INITIAL THERAPY**

**SUBSEQUENT THERAPY<sup>g</sup>**



### Regimen Dosing (SBA-D 4 of 7)

<sup>a</sup> Many of the regimens recommended in these guidelines are extrapolated from data for colorectal cancer.

<sup>b</sup> For infection risk, monitoring, and prophylaxis recommendations for targeted therapies, see INF-A in the [NCCN Guidelines for Prevention and Treatment of Cancer-Related Infections](#).

<sup>e</sup> See [NCCN Guidelines for Management of Immunotherapy-Related Toxicities](#).

<sup>f</sup> If no previous treatment with a checkpoint inhibitor.

<sup>g</sup> Larotrectinib or entrectinib are treatment options for patients with metastatic SBA that is *NTRK* gene fusion positive.

**Note: All recommendations are category 2A unless otherwise indicated.**  
**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**



**ADJUVANT THERAPY REGIMENS<sup>a,h</sup>**

- **mFOLFOX6<sup>1-3</sup>**
  - ▶ Oxaliplatin 85 mg/m<sup>2</sup> IV, day 1<sup>i</sup>
  - ▶ Leucovorin 400 mg/m<sup>2</sup> IV, day 1<sup>j</sup>
  - ▶ 5-FU 400 mg/m<sup>2</sup> IV bolus on day 1, then 1200 mg/m<sup>2</sup>/day x 2 days (total 2400 mg/m<sup>2</sup> over 46–48 hours) continuous infusion
  - ▶ Repeat every 2 weeks
- **5-FU/leucovorin<sup>1-3</sup>**
  - ▶ Leucovorin 500 mg/m<sup>2</sup> given as a 2-hour infusion and repeated weekly x 6<sup>j</sup>
  - ▶ 5-FU 500 mg/m<sup>2</sup> given bolus 1 hour after the start of leucovorin and repeated weekly x 6
  - ▶ Every 8 weeks for 4 cycles
- **Simplified biweekly infusional 5-FU/LV (sLV5FU2)<sup>1-3</sup>**
  - ▶ Leucovorin 400 mg/m<sup>2</sup> IV day 1,<sup>j</sup>
  - ▶ followed by 5-FU bolus 400 mg/m<sup>2</sup> and then 1200 mg/m<sup>2</sup>/day x 2 days (total 2400 mg/m<sup>2</sup> over 46–48 hours) continuous infusion
  - ▶ Repeat every 2 weeks
- **Capecitabine<sup>1-3</sup>**
  - ▶ Capecitabine 1000–1250 mg/m<sup>2</sup> twice daily for 14 days every 3 weeks x 24 weeks
- **Capecitabine + RT<sup>4</sup>**
  - ▶ Capecitabine 825 mg/m<sup>2</sup> PO twice daily on days 1–7 for 5 weeks with RT
- **5-FU with RT<sup>4</sup>**
  - ▶ 5-FU 225 mg/m<sup>2</sup> over 24 hours daily on days 1–7 for 5 weeks with RT
- **CAPEOX<sup>1-3</sup>**
  - ▶ Oxaliplatin 130 mg/m<sup>2</sup> IV day 1<sup>i</sup>
  - ▶ Capecitabine 1000 mg/m<sup>2</sup> twice daily for 14 days every 3 weeks x 24 weeks<sup>k</sup>

**See References (SBA-D 7 of 7)**

<sup>a</sup> Many of the regimens recommended in these guidelines are extrapolated from data for colorectal cancer.

<sup>h</sup> The role for adjuvant chemotherapy remains controversial due to conflicting results across a number of retrospective analyses. Participation in clinical trials is strongly recommended.

<sup>i</sup> Oxaliplatin may be given either over 2 hours, or may be infused over a shorter time at a rate of 1 mg/m<sup>2</sup>/min. Leucovorin infusion should match infusion time of oxaliplatin. Cerek A, Park V, Yaeger R, et al. Faster FOLFOX: oxaliplatin can be safely infused at a rate of 1 mg/m<sup>2</sup>/min. J Oncol Pract 2016;12:e548-553.

<sup>j</sup> Leucovorin 400 mg/m<sup>2</sup> is the equivalent of levoleucovorin 200 mg/m<sup>2</sup>.

<sup>k</sup> The majority of safety and efficacy data for this regimen have been developed in Europe, where a capecitabine starting dose of 1000 mg/m<sup>2</sup> twice daily for 14 days, repeated every 21 days, is standard. Evidence suggests that North American patients may experience greater toxicity with capecitabine (as well as with other fluoropyrimidines) than European patients, and may require a lower dose of capecitabine.

**Note: All recommendations are category 2A unless otherwise indicated.**

**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**

**ADVANCED OR METASTATIC THERAPY REGIMENS<sup>a</sup>**

- **FOLFOX**
  - ▶ **mFOLFOX<sup>6-7</sup>**
    - ◇ Oxaliplatin 85 mg/m<sup>2</sup> IV day 1<sup>i</sup>
    - ◇ Leucovorin 400 mg/m<sup>2</sup> IV day 1<sup>j</sup>
    - ◇ 5-FU 400 mg/m<sup>2</sup> IV bolus on day 1, then 1200 mg/m<sup>2</sup>/day x 2 days (total 2400 mg/m<sup>2</sup> over 46–48 hours) IV continuous infusion
    - ◇ Repeat every 2 weeks
  - ▶ **mFOLFOX<sup>7-8</sup>**
    - ◇ Oxaliplatin 85 mg/m<sup>2</sup> IV day 1<sup>i</sup>
    - ◇ Leucovorin 400 mg/m<sup>2</sup> IV day 1<sup>j</sup>
    - ◇ 5-FU 1200 mg/m<sup>2</sup>/day x 2 days (total 2400 mg/m<sup>2</sup> over 46–48 hours) IV continuous infusion
    - ◇ Repeat every 2 weeks
- **FOLFOX + bevacizumab<sup>9,10</sup>**
  - ▶ Bevacizumab 5 mg/kg IV, day 1<sup>l</sup>
  - ▶ Repeat every 2 weeks
- **CAPEOX<sup>11</sup>**
  - ▶ Oxaliplatin 130 mg/m<sup>2</sup> IV day 1<sup>i</sup>
  - ▶ Capecitabine 1000 mg/m<sup>2</sup> twice daily PO for 14 days<sup>k</sup>
  - ▶ Repeat every 3 weeks
- **CAPEOX + bevacizumab<sup>9,10,12</sup>**
  - ▶ Oxaliplatin 130 mg/m<sup>2</sup> IV day 1<sup>i</sup>
  - ▶ Capecitabine 1000 mg/m<sup>2</sup> PO twice daily for 14 days<sup>k</sup>
  - ▶ Bevacizumab 7.5 mg/kg IV day 1<sup>l</sup>
  - ▶ Repeat every 3 weeks
- **FOLFIRI<sup>13</sup>**
  - ▶ Irinotecan 180 mg/m<sup>2</sup> IV over 30–90 minutes, day 1
  - ▶ Leucovorin 400 mg/m<sup>2</sup> IV infusion to match duration of irinotecan infusion, day 1<sup>j</sup>
  - ▶ 5-FU 400 mg/m<sup>2</sup> IV bolus day 1, then 1200 mg/m<sup>2</sup>/day x 2 days (total 2400 mg/m<sup>2</sup> over 46–48 hours) continuous infusion
  - ▶ Repeat every 2 weeks
- **FOLFOXIRI<sup>14,15</sup>**
  - ▶ Irinotecan 165 mg/m<sup>2</sup> IV day 1
  - ▶ Oxaliplatin 85 mg/m<sup>2</sup> IV day 1<sup>i</sup>
  - ▶ Leucovorin 400 mg/m<sup>2</sup> day 1<sup>j</sup>
  - ▶ Fluorouracil 1200 mg/m<sup>2</sup>/day x 2 days (total 2400 mg/m<sup>2</sup> over 48 hours) continuous infusion starting on day 1
  - ▶ Repeat every 2 weeks
  - ▶ The dose used in European studies was 3200 mg/m<sup>2</sup>. U.S. patients have been shown to have poorer tolerance for 5-FU. The dose listed above is recommended for U.S. patients.
- **FOLFOXIRI + bevacizumab<sup>9,10</sup>**
  - ▶ Bevacizumab 5 mg/kg IV, day 1<sup>l</sup>
  - ▶ Repeat every 2 weeks

[See Additional Regimens \(SBA-D 5 of 7\)](#)  
[See References \(SBA-D 7 of 7\)](#)

<sup>a</sup> Many of the regimens recommended in these guidelines are extrapolated from data for colorectal cancer.

<sup>i</sup> Oxaliplatin may be given either over 2 hours, or may be infused over a shorter time at a rate of 1 mg/m<sup>2</sup>/min. Leucovorin infusion should match infusion time of oxaliplatin. Cerek A, Park V, Yaeger R, et al. Faster FOLFOX: oxaliplatin can be safely infused at a rate of 1 mg/m<sup>2</sup>/min. J Oncol Pract 2016;12:e548-553.

<sup>j</sup> Leucovorin 400 mg/m<sup>2</sup> is the equivalent of levoleucovorin 200 mg/m<sup>2</sup>.

<sup>k</sup> The majority of safety and efficacy data for this regimen have been developed in Europe, where a capecitabine starting dose of 1000 mg/m<sup>2</sup> twice daily for 14 days, repeated every 21 days, is standard. Evidence suggests that North American patients may experience greater toxicity with capecitabine (as well as with other fluoropyrimidines) than European patients, and may require a lower dose of capecitabine.

<sup>l</sup> Bevacizumab may be safely given at a rate of 0.5 mg/kg/min (5 mg/kg over 10 minutes and 7.5 mg/kg over 15 minutes).

**Note: All recommendations are category 2A unless otherwise indicated.**

**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**

**ADVANCED OR METASTATIC THERAPY REGIMENS<sup>a</sup>**

- **Capecitabine + RT<sup>4</sup>**
  - ▶ Capecitabine 825 mg/m<sup>2</sup> PO twice daily on Days 1–7 for 5 weeks with RT
- **5-FU with RT<sup>4</sup>**
  - ▶ 5-FU 225 mg/m<sup>2</sup> over 24 hours daily on days 1–7 for 5 weeks with RT
- **5-FU/leucovorin**
  - ▶ **Roswell Park regimen<sup>16</sup>**
    - ◊ Leucovorin 500 mg/m<sup>2</sup> IV over 2 hours, days 1, 8, 15, 22, 29, and 36<sup>9</sup>
    - ◊ 5-FU 500 mg/m<sup>2</sup> IV bolus 1 hour after start of leucovorin, days 1, 8, 15, 22, 29, and 36
    - ◊ Repeat every 8 weeks
  - ▶ **Simplified biweekly infusional 5-FU/LV (sLV5FU2)<sup>17</sup>**
    - ◊ Leucovorin 400 mg/m<sup>2</sup> IV over 2 hours on day 1,<sup>j</sup>
    - ◊ followed by 5-FU bolus 400 mg/m<sup>2</sup> and then 1200 mg/m<sup>2</sup>/day x 2 days (total 2400 mg/m<sup>2</sup> over 46–48 hours) continuous infusion
    - ◊ Repeat every 2 weeks
  - ▶ **Weekly<sup>18</sup>**
    - ◊ Leucovorin 20 mg/m<sup>2</sup> IV over 2 hours on day 1<sup>j</sup>
    - ◊ 5-FU 500 mg/m<sup>2</sup> IV bolus injection 1 hour after the start of leucovorin
    - ◊ Repeat every week
    - or
    - ◊ 5-FU 2600 mg/m<sup>2</sup> by 24-hour infusion on day 1
    - ◊ Leucovorin 500 mg/m<sup>2</sup> over 2 hours on day 1<sup>j</sup>
    - ◊ Repeat every week
- **Capecitabine<sup>19</sup>**
  - ▶ Capecitabine 850–1250 mg/m<sup>2</sup> PO twice daily, for 14 days
  - ▶ Repeat every 3 weeks
- **Irinotecan<sup>20,21</sup>**
  - ▶ Irinotecan 125 mg/m<sup>2</sup> IV over 30–90 minutes, days 1 and 8
  - ▶ Repeat every 3 weeks
  - or
  - ▶ Irinotecan 180 mg/m<sup>2</sup> IV over 30–90 minutes, day 1
  - ▶ Repeat every 2 weeks or
  - or
  - ▶ Irinotecan 300–350 mg/m<sup>2</sup> IV over 30–90 minutes, day 1
  - ▶ Repeat every 3 weeks
- **Pembrolizumab (dMMR/MSI-H only)<sup>22</sup>**
  - ▶ Pembrolizumab 2 mg/kg IV every 3 weeks
  - or
  - ▶ Pembrolizumab 200 mg IV every 3 weeks
  - or
  - ▶ Pembrolizumab 400 mg IV every 6 weeks
- **Nivolumab (dMMR/MSI-H only)<sup>23</sup>**
  - ▶ Nivolumab 3 mg/kg every 2 weeks
  - or
  - ▶ Nivolumab 240 mg IV every 2 weeks
  - or
  - ▶ Nivolumab 480 mg IV every 4 weeks
- **Ipilimumab + nivolumab (dMMR/MSI-H only)<sup>24</sup>**
  - ▶ Nivolumab 3 mg/kg (30-minute IV infusion)
  - ▶ Ipilimumab 1 mg/kg (30-minute IV infusion)
  - ▶ Once every 3 weeks for four doses, then
  - ▶ Nivolumab 3 mg/kg IV or nivolumab 240 mg IV every 2 weeks
- **Dostarlimab-gxly<sup>25</sup> (dMMR/MSI-H only)**
  - ▶ Dostarlimab-gxly 500 mg IV every 3 weeks for 4 doses followed by 1,000 mg IV every 6 weeks

<sup>a</sup> Many of the regimens recommended in these guidelines are extrapolated from data for colorectal cancer.

<sup>j</sup> Leucovorin 400 mg/m<sup>2</sup> is the equivalent of levoleucovorin 200 mg/m<sup>2</sup>.

[See Additional Regimens \(SBA-D 6 of 7\)](#)

[See References \(SBA-D 7 of 7\)](#)

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**ADVANCED OR METASTATIC THERAPY REGIMENS<sup>a</sup>**

- **Albumin-bound paclitaxel<sup>26</sup>**
  - ▶ Albumin-bound paclitaxel 220–260 mg/m<sup>2</sup> IV every 21 days
- **Docetaxel<sup>27</sup>**
  - ▶ Docetaxel 75–100 mg/m<sup>2</sup> IV on day 1 every 21 days
- **Paclitaxel<sup>27</sup>**
  - ▶ Paclitaxel 135–250 mg/m<sup>2</sup> IV on day 1 every 21 days
  - or
  - ▶ Paclitaxel 80 mg/m<sup>2</sup> IV weekly
  - or
  - ▶ Paclitaxel 80 mg/m<sup>2</sup> IV on days 1, 8, 15 every 28 days
- **Gemcitabine + albumin-bound paclitaxel<sup>27</sup>**
  - ▶ Albumin-bound paclitaxel 125 mg/m<sup>2</sup> IV on Days 1, 8, 15
  - ▶ Gemcitabine 1000 mg/m<sup>2</sup> IV on Days 1, 8, 15
  - ▶ Every 28 days
- **Gemcitabine + docetaxel<sup>27</sup>**
  - ▶ Gemcitabine 1000 mg/m<sup>2</sup> IV on days 1 and 8
  - ▶ Docetaxel 75 mg/m<sup>2</sup> IV on Day 8
  - ▶ Every 21 days
- **Gemcitabine + paclitaxel<sup>27</sup>**
  - ▶ Gemcitabine 1000 mg/m<sup>2</sup> IV on days 1, 8, 15
  - ▶ Paclitaxel 110 mg/m<sup>2</sup> on days 1, 8, 15
  - ▶ Every 28 days
- **Carboplatin + paclitaxel<sup>27</sup>**
  - ▶ Paclitaxel 175 mg/m<sup>2</sup> IV on Day 1
  - ▶ Carboplatin AUC 5 IV on Day 1
  - ▶ Every 21 days
- **Gemcitabine, docetaxel, and capecitabine (GTX)<sup>27</sup>**
  - ▶ Gemcitabine 750 mg/m<sup>2</sup> IV at a rate of 10 mg/m<sup>2</sup>/min on days 4 and 11
  - ▶ Docetaxel 30 mg/m<sup>2</sup> IV on days 4 and 11
  - ▶ Capecitabine 750 mg/m<sup>2</sup> PO twice daily for 14 days
  - ▶ Every 21 days for 2–6 cycles
- **Larotrectinib<sup>28</sup> (*NTRK* gene fusion positive)**
  - ▶ 100 mg PO twice daily
- **Entrectinib<sup>29</sup> (*NTRK* gene fusion positive)**
  - ▶ 600 mg PO once daily

See References (SBA-D 7 of 7)

<sup>a</sup> Many of the regimens recommended in these guidelines are extrapolated from data for colorectal cancer.**Note: All recommendations are category 2A unless otherwise indicated.****Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**

### PRINCIPLES OF SYSTEMIC THERAPY - REFERENCES

- 1 Evans TRJ, Aparicio T, Malicot KL, et al. GLOBAL BALLAD: An International Rare Cancers Initiative trial to evaluate the potential benefit of adjuvant chemotherapy for small bowel adenocarcinoma (IRCI 002) [abstract]. *J Clin Oncol* 2016;34:TPS4154-TPS4154.
- 2 Ecker BL, McMillan MT, Datta J, et al. Efficacy of adjuvant chemotherapy for small bowel adenocarcinoma: A propensity score-matched analysis. *Cancer* 2016;122:693-701.
- 3 Overman MJ, Kopetz S, Lin E, et al. Is there a role for adjuvant therapy in resected adenocarcinoma of the small intestine. *Acta Oncol* 2010;49:474-479.
- 4 Allegra CJ, Yothers G, O'Connell MJ, et al. Neoadjuvant 5-FU or capecitabine plus radiation with or without oxaliplatin in rectal cancer patients: A phase III randomized clinical trial. *J Natl Cancer Inst* 2015;107:djv248.
- 5 Horimatsu T, Nakayama N, Moriwaki T, et al. A phase II study of 5-fluorouracil/L-leucovorin/oxaliplatin (mFOLFOX6) in Japanese patients with metastatic or unresectable small bowel adenocarcinoma. *Int J Clin Oncol* 2017;22:905-912.
- 6 Tsushima T, Taguri M, Honma Y, et al. Multicenter retrospective study of 132 patients with unresectable small bowel adenocarcinoma treated with chemotherapy. *Oncologist* 2012;17:1163-1170.
- 7 Zaanan A, Costes L, Gauthier M, et al. Chemotherapy of advanced small-bowel adenocarcinoma: a multicenter AGEO study. *Ann Oncol* 2010;21:1786-1793.
- 8 Xiang XJ, Liu YW, Zhang L, et al. A phase II study of modified FOLFOX as first-line chemotherapy in advanced small bowel adenocarcinoma. *Anticancer Drugs* 2012;23:561-566.
- 9 Aydin D, Sendur MA, Kefeli U, et al. Evaluation of bevacizumab in advanced small bowel adenocarcinoma. *Clin Colorectal Cancer* 2017;16:78-83.
- 10 Takayoshi K, Kusaba H, Uenomachi M, et al. Suggestion of added value by bevacizumab to chemotherapy in patients with unresectable or recurrent small bowel cancer. *Cancer Chemother Pharmacol* 2017;80:333-342.
- 11 Overman MJ, Varadhachary GR, Kopetz S, et al. Phase II study of capecitabine and oxaliplatin for advanced adenocarcinoma of the small bowel and ampulla of Vater. *J Clin Oncol* 2009;27:2598-2603.
- 12 Gulhati P, Raghav K, Shroff RT, et al. Bevacizumab combined with capecitabine and oxaliplatin in patients with advanced adenocarcinoma of the small bowel or ampulla of vater: A single-center, open-label, phase 2 study. *Cancer* 2017;123:1011-1017.
- 13 Zaanan A, Gauthier M, Malka D, et al. Second-line chemotherapy with fluorouracil, leucovorin, and irinotecan (FOLFIRI regimen) in patients with advanced small bowel adenocarcinoma after failure of first-line platinum-based chemotherapy: a multicenter AGEO study. *Cancer* 2011;117:1422-1428.
- 14 Masi G, Allegrini G, Cupini S, et al. First-line treatment of metastatic colorectal cancer with irinotecan, oxaliplatin, and 5-fluorouracil/leucovorin (FOLFOXIRI): results of a phase II study with a simplified biweekly schedule. *Ann Oncol* 2004;15:1766-1772.
- 15 McWilliams RR, Foster NR, Mahoney MR, et al. North Central Cancer Treatment Group N0543 (Alliance): A phase 2 trial of pharmacogenetic-based dosing of irinotecan, oxaliplatin, and capecitabine as first-line therapy for patients with advanced small bowel adenocarcinoma. *Cancer* 2017;123:3494-3501.
- 16 Wolmark N, Rockette H, Fisher B, et al. The benefit of leucovorin-modulated fluorouracil as postoperative adjuvant therapy for primary colon cancer: results from National Surgical Adjuvant Breast and Bowel Project protocol C-03. *J Clin Oncol* 1993;11:1879-1887.
- 17 Andre T, Louvet C, Maindault-Goebel F, et al. CPT-11 (irinotecan) addition to bimonthly, high-dose leucovorin and bolus and continuous-infusion 5-fluorouracil (FOLFIRI) for pretreated metastatic colorectal cancer. *Eur J Cancer* 1999;35(9):1343-7.
- 18 Jäger E, Heike M, Bernhard H, et al. Weekly high-dose leucovorin versus low-dose leucovorin combined with fluorouracil in advanced colorectal cancer: results of a randomized multicenter trial. Study Group for Palliative Treatment of Metastatic Cancer Study Protocol 1. *J Clin Oncol* 1996;14:2274-2279.
- 19 Saltz LB, Clarke S, Diaz-Rubio E, et al. Bevacizumab in combination with oxaliplatin-based chemotherapy as first-line therapy in metastatic colorectal cancer: a randomized phase III study. *J Clin Oncol* 2008;26:2013-2019.
- 20 Polyzos A, Kouraklis G, Giannopoulos A, et al. Irinotecan as salvage chemotherapy for advanced small bowel adenocarcinoma: a series of three patients. *J Chemother* 2003;15:503-506.
- 21 Suenaga M, Mizunuma N, Chin K, et al. Chemotherapy for small-bowel Adenocarcinoma at a single institution. *Surg Today* 2009;39:27-31.
- 22 Le DT, Uram JN, Wang H, et al. PD-1 Blockade in tumors with mismatch-repair deficiency. *N Engl J Med* 2015;372:2509-2520.
- 23 Overman MJ, McDermott R, Leach JL, et al. Nivolumab in patients with metastatic DNA mismatch repair-deficient or microsatellite instability-high colorectal cancer (CheckMate 142): an open-label, multicentre, phase 2 study. *Lancet Oncol* 2017;18:1182-1191.
- 24 Overman MJ, Lonardi S, Wong KYM, et al. Durable clinical benefit with nivolumab plus ipilimumab in DNA mismatch repair-deficient/microsatellite instability-high metastatic colorectal cancer. *J Clin Oncol* 2018;36:773-779.
- 25 Berton D, Banerjee SN, Curigliano G, et al. Antitumor activity of dostarlimab in patients with mismatch repair-deficient/microsatellite instability-high tumors: A combined analysis of two cohorts in the GARNET study. *J Clin Oncol* 2021;39:15\_suppl, 2564-2564.
- 26 Overman MJ, Adam L, Raghav K, et al. Phase II study of nab-paclitaxel in refractory small bowel adenocarcinoma and CpG island methylator phenotype (CIMP)-high colorectal cancer. *Ann Oncol* 2018;29:139-144.
- 27 Aldrich JD, Raghav KPS, Varadhachary GR, Wolff RA, Overman MJ. Retrospective analysis of taxane-based therapy in small bowel adenocarcinoma. *Oncologist* 2019;24:e384-e386.
- 28 Drilon A, Laetsch TW, Kummar S, et al. Efficacy of larotrectinib in *TRK* fusion positive cancers in adults and children. *N Engl J Med* 2018;378:731-739.
- 29 Doebele RC, Drilon A, Paz-Ares L, et al. Entrectinib in patients with advanced or metastatic NTRK fusion-positive solid tumours: integrated analysis of three phase 1-2 trials. *Lancet Oncol* 2020;21:271-282.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PRINCIPLES OF RADIATION THERAPY****Duodenum:**

- Database analysis suggests no survival benefit from the addition of adjuvant chemoradiation versus chemotherapy alone in patients with surgically resected duodenal adenocarcinoma.<sup>1</sup> A separate retrospective study showed mixed results regarding the efficacy of either preoperative or postoperative chemoradiation for the management of locally advanced or margin-positive duodenal adenocarcinomas.<sup>2</sup> Therefore, chemoradiation should be considered only in highly select patients.
- Preoperative chemoradiation should be considered in patients who remain unresectable following a course of induction chemotherapy.
- Patients should be evaluated by multidisciplinary teams at high-volume centers in cases where either preoperative or postoperative radiation therapy (RT) is being considered.
- Treatment Information:
  - ▶ Fluoropyrimidine-based chemotherapy should be delivered concurrently with RT.
  - ▶ Treatment can be delivered using 3D conformal RT. When appropriate, advanced treatment planning, such as intensity-modulated RT (IMRT), should be considered to limit toxicity to adjacent normal organs.
  - ▶ Image-guided RT (IGRT) with kilovoltage (kV) imaging, MR-guided imaging, and cone beam CT imaging should be routinely used during the course of treatment with IMRT.
  - ▶ Target Volumes:
    - ◊ The primary site and regional lymph node basins should be included in the RT fields.
  - ▶ RT Dosing:
    - ◊ Doses of 45–54 Gy in 1.8–2 Gy daily fractions should be used based on tolerance limits of adjacent normal tissues.
    - ◊ Adjacent small bowel dose should be limited to 45 Gy, if possible.

**Jejunum/Ileum:**

- RT is not generally indicated for lesions arising in these sites. Any consideration for such therapy must be made on a highly selected basis by a multidisciplinary team.

<sup>1</sup>Ecker BL, McMillan MT, Datta J, et al. Adjuvant chemotherapy versus chemoradiotherapy in the management of patients with surgically resected duodenal adenocarcinoma: A propensity score-matched analysis of a nationwide clinical oncology database. *Cancer* 2017;123:967-976.

<sup>2</sup>Kelsey CR, Nelson JW, Willett CG, et al. Duodenal adenocarcinoma: patterns of failure after resection and the role of chemoradiotherapy. *Int J Radiat Oncol Biol Phys* 2007;69:1436-1441.

**Note:** All recommendations are category 2A unless otherwise indicated.

**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



### PRINCIPLES OF SURVIVORSHIP

- **SBA surveillance, [see SBA-2](#)**
- **Survivorship Care Planning**
  - ▶ **Should have defined roles for the oncologist and the primary care provider following curative-intent therapy**
  - ▶ **Overall summary, including dates, of all diagnostic and surgical procedures, chemotherapy received (including drug and number of cycles), and radiation (including dose) should be provided to the patient at end of treatment**
  - ▶ **Description of long-term toxicities and when they may resolve, as well as potential delayed toxicity**
  - ▶ **Surveillance recommendations and schedule**
  - ▶ **Health behavior recommendations**
- **Management of Late/Long-Term Sequelae of Treatment**
  - ▶ **For issues related to distress, pain, neuropathy, fatigue, or sexual dysfunction, [see NCCN Guidelines for Survivorship](#)**
- **Oxaliplatin-Related Neuropathy**
  - ▶ **Only consider duloxetine, pregabalin, or gabapentin in cases where pain is present, as these are ineffective in treating numbness or tingling associated with the long-term use of oxaliplatin**
  - ▶ **Consider non-pharmacologic interventions, including balanced physical activity, acupuncture, heat or ice, or other methods**
- **Lifestyle Modifications**
  - ▶ **Avoid gluten in patients with confirmed celiac disease**
  - ▶ **Eat a plant-based diet**
  - ▶ **Maintain healthy body weight throughout life**
  - ▶ **Engage in regular physical activity**
  - ▶ **Undergo all age- and gender-appropriate preventive health and cancer screenings at recommended intervals with primary care provider**
  - ▶ **Avoid the use of alcohol or tobacco**
- **Crohn's Disease**
  - ▶ **Patients with Crohn's disease and history of SBA remain at elevated risk for developing further SBAs. Surveillance screening should be considered for these individuals.<sup>1,2</sup>**

<sup>1</sup> Grolleau C, Pote NM, Guedj NS, et al. Small bowel adenocarcinoma complicating Crohn's disease: a single-centre experience emphasizing the importance of screening for dysplasia. *Virchows Arch* 2017;471:611-617.

<sup>2</sup> Cahill C, Gordon PH, Petrucci A, Boutros M. Small bowel adenocarcinoma and Crohn's disease: any further ahead than 50 years ago? *World J Gastroenterol* 2014;20:11486-11495.

**Note: All recommendations are category 2A unless otherwise indicated.**

**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**

**American Joint Committee on Cancer (AJCC) TNM Staging Classification for Small Intestine Adenocarcinoma 8th ed., 2017****Table 1. Definitions for T, N, M**

<b>T</b>	<b>Primary Tumor</b>
<b>TX</b>	Primary tumor cannot be assessed
<b>T0</b>	No evidence of primary tumor
<b>Tis</b>	High-grade dysplasia/carcinoma <i>in situ</i>
<b>T1</b>	Tumor invades the lamina propria or submucosa
T1a	Tumor invades the lamina propria
T1b	Tumor invades the submucosa
<b>T2</b>	Tumor invades the muscularis propria
<b>T3</b>	Tumor invades through the muscularis propria into the subserosa, or extends into nonperitonealized perimuscular tissue (mesentery or retroperitoneum) without serosal penetration
<b>T4</b>	Tumor perforates the visceral peritoneum or directly invades other organs or structures (e.g., other loops of small intestine, mesentery of adjacent loops of bowel, and abdominal wall by way of serosa; for duodenum only, invasion of pancreas or bile duct)
<b>N</b>	<b>Regional Lymph Nodes</b>
<b>NX</b>	Regional lymph nodes cannot be assessed
<b>N0</b>	No regional lymph node metastasis
<b>N1</b>	Metastasis in one or two regional lymph nodes
<b>N2</b>	Metastasis in three or more regional lymph nodes
<b>M</b>	<b>Distance Metastasis</b>
<b>M0</b>	No distant metastasis
<b>M1</b>	Distant metastasis present

**Table 2. AJCC Prognostic Stage Groups**

	<b>T</b>	<b>N</b>	<b>M</b>
<b>Stage 0</b>	Tis	N0	M0
<b>Stage I</b>	T1-2	N0	M0
<b>Stage IIA</b>	T3	N0	M0
<b>Stage IIB</b>	T4	N0	M0
<b>Stage IIIA</b>	Any T	N1	M0
<b>Stage IIIB</b>	Any T	N2	M0
<b>Stage IV</b>	Any T	Any N	M1

Used with the permission of the American College of Surgeons, Chicago Illinois. The original source for this information is the AJCC Cancer Staging Manual, Eighth Edition (2017) published by Springer International Publishing.





### NCCN Categories of Evidence and Consensus

<b>Category 1</b>	Based upon high-level evidence, there is uniform NCCN consensus that the intervention is appropriate.
<b>Category 2A</b>	Based upon lower-level evidence, there is uniform NCCN consensus that the intervention is appropriate.
<b>Category 2B</b>	Based upon lower-level evidence, there is NCCN consensus that the intervention is appropriate.
<b>Category 3</b>	Based upon any level of evidence, there is major NCCN disagreement that the intervention is appropriate.

All recommendations are category 2A unless otherwise indicated.



# NCCN Guidelines Version 2.2021

## Small Bowel Adenocarcinoma

### Discussion

This discussion corresponds to the NCCN Guidelines for Small Bowel Adenocarcinoma. Last updated July 30, 2019

### Table of Contents

**Overview** .....MS-2

**Literature Search Criteria and Guidelines Update Methodology** .MS-2

**Risk Factors for Small Bowel Adenocarcinoma**.....MS-3

    Lifestyle Factors.....MS-3

    Inflammatory Bowel Disease and Celiac Disease .....MS-3

    Familial Syndromes .....MS-3

**Clinical Presentation and Workup**.....MS-4

    Imaging and Endoscopy.....MS-5

**Pathology and Staging**.....MS-5

    Lymph Node Evaluation .....MS-6

**Treatment of Stage I-III Small Bowel Adenocarcinoma**.....MS-6

    Surgical Management of Localized Resectable Disease .....MS-6

    Adjuvant Therapy.....MS-7

    Primary Treatment of Unresectable Disease.....MS-8

**Treatment of Distant Metastatic (Stage IV) SBA**.....MS-9

    Metastasectomy.....MS-9

    Peritoneal Carcinomatosis .....MS-9


    Systemic Therapy for Metastatic Disease .....MS-10

**Posttreatment Surveillance** ..... MS-13

**Survivorship** ..... MS-14

**Summary**..... MS-14

**References**..... MS-15





# NCCN Guidelines Version 2.2021

## Small Bowel Adenocarcinoma

### Overview

In 2019, an estimated 10,590 new cases of small bowel cancer will occur and 1590 patients will die of this disease.<sup>1</sup> Compared to cancers of other organs in the gastrointestinal tract, small bowel cancers are relatively rare, accounting for only about 3% of cancers occurring in this organ system.<sup>1</sup> Small bowel cancers affect men and women relatively equally, with an incidence of 2.6 per 100,000 for men and 2.0 per 100,000 for women.<sup>2</sup> The median age at diagnosis is 66 years. The incidence of small bowel cancers is increasing, with an annual percent increase of 1.8 between 2006 and 2015. This trend is in contrast to other gastrointestinal malignancies, including esophageal, gastric, colon, and rectum, which decreased in incidence across the same timeframe.<sup>2</sup> The four most common histologies of cancers originating in the small bowel are adenocarcinomas, neuroendocrine tumors, gastrointestinal stromal tumors (GIST), and lymphomas.<sup>3,4</sup> The treatment recommendations in this guideline only refer to small bowel adenocarcinoma (SBA), which comprise an estimated 30% to 40% incidence of small intestinal cancer diagnoses.<sup>4</sup> Due to the rarity of this disease, there are very few established guidelines for management of SBA. In 2018, a French intergroup published the first clinical practice guidelines for SBA.<sup>5</sup> These NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Small Bowel Adenocarcinoma are the second.

This Discussion summarizes the NCCN Guidelines for Small Bowel Adenocarcinoma. These guidelines begin with the clinical presentation of the patient to the primary care physician or gastroenterologist and address diagnosis, pathologic staging, surgical management, perioperative treatment, patient surveillance, management of recurrent and metastatic disease, and survivorship. When reviewing these guidelines, clinicians should be aware of several things. First, these guidelines adhere to the TNM (tumor, node, metastases) staging system (Table 1 in the algorithm).<sup>6</sup> Furthermore, all recommendations are classified as category

2A except where noted in the text or algorithm. Although the guidelines are believed to represent the optimal treatment strategy, ***participation in a clinical trial is especially encouraged for patients with SBA based on the dearth of clinical trial data on which to base treatment decisions for this disease.***

### Literature Search Criteria and Guidelines Update Methodology

Prior to the development of the NCCN Guidelines for Small Bowel Adenocarcinoma, an electronic search of the PubMed database was performed to obtain key literature in the field of small bowel cancer, using the following search terms: (small bowel cancer) OR (small intestine cancer) OR (jejunum cancer) OR (duodenum cancer) OR (ileum cancer). The PubMed database was chosen because it remains the most widely used resource for medical literature and indexes only peer-reviewed biomedical literature.<sup>7</sup> The search results were narrowed by selecting studies in humans published in English. Results were confined to the following article types: Clinical Trial; Multicenter Study; Practice Guideline; Randomized Controlled Trial; Meta-Analysis; Systematic Reviews; and Validation Studies.

The data from key PubMed articles and articles from additional sources deemed as relevant to these Guidelines and discussed by the panel have been included in this version of the Discussion section (eg, e-publications ahead of print, meeting abstracts). Recommendations for which high-level evidence is lacking are based on the panel's review of lower-level evidence and expert opinion.

The complete details of the Development and Update of the NCCN Guidelines are available on the NCCN website at [www.NCCN.org](http://www.NCCN.org).



# NCCN Guidelines Version 2.2021

## Small Bowel Adenocarcinoma

### Risk Factors for Small Bowel Adenocarcinoma

Risk factors for SBA are similar to those for colorectal cancer (CRC), including lifestyle factors, inflammatory bowel disease (IBD), and certain familial syndromes such as Lynch syndrome (also known as hereditary nonpolyposis colorectal cancer [HNPCC]), Peutz-Jeghers syndrome (PJS), and familial adenomatous polyposis (FAP). Therefore, it is recommended that all patients with small bowel cancer be queried regarding their family history and considered for risk assessment, as detailed in the [NCCN Guidelines for Colorectal Cancer Screening](#).

### Lifestyle Factors

Although data on the role of lifestyle factors in relation to the risk of developing SBA are very limited due to low incidence of disease, lifestyle factors that have been reported as raising the risk of SBA generally agree with known risk factors for CRC. A systematic review of the literature has reported that high levels of alcohol consumption, smoking, and dietary factors, including low intake of fiber and high intake of red/processed meat and sugary drinks, may increase the risk of SBA.<sup>8</sup> Additionally, the results of a pooled analysis of more than 500,000 subjects in the Asia Cohort Consortium reported that elevated body mass index (BMI) and high alcohol consumption were associated with a non-significant trend towards an increased risk of SBA, although this analysis did not identify smoking as a risk factor.<sup>9</sup>

### Inflammatory Bowel Disease and Celiac Disease

It is well recognized that individuals with IBD (ulcerative colitis or Crohn's disease) are at increased risk for CRC. Several studies have also reported an increased risk of distal SBA in patients with IBD.<sup>10-13</sup> The results of a retrospective, multicenter observational cohort study of 9100 patients with IBD found that the relative risk of small bowel cancer was 3.70 (95% CI, 1.23–11.13) for patients with IBD. The rate of death and cancer remission did not differ between patients who maintained treatment for IBD

compared to those who stopped IBD treatment.<sup>10</sup> Additionally, although the data are mostly limited to case studies and literature reviews, cases of SBA have been reported in patients with celiac disease suggesting a possible link between these conditions.<sup>14-16</sup> The association with celiac disease is poorly understood and a distinct difference from CRC, for which celiac disease is not a risk factor.

### Familial Syndromes

Due to the relative rarity of SBA, and the disease's association with several genetic syndromes, the NCCN Panel recommends that all patients with a personal history of SBA should be counseled for familial malignancies and considered for risk assessment of various genetic syndromes, including Lynch syndrome, PJS, FAP, and other polypoid mutations. A brief description of some of these syndromes is included below. See the [NCCN Guidelines for Genetic/Familial High-Risk Assessment: Colorectal](#) for more information.

#### **Familial Adenomatous Polyposis**

FAP is an autosomal dominant condition characterized by a germline mutation in the *APC* gene, located on chromosome 5q21.<sup>17,18</sup> Patients with FAP develop large numbers of adenomatous polyps in the large bowel, beginning as early as adolescence, and most patients with classic FAP will develop polyps by the age of 25. Patients with attenuated FAP due to a germline *MUTYH* mutation develop fewer numbers of polyps, generally at a later age than those with classic FAP.<sup>17,18</sup> While the incidence of SBA in patients with FAP has not been well established, the lifetime risk has been estimated as 3% to 5%.<sup>19</sup> The duodenum and periampullary region are the most common locations for SBA in patients with FAP.<sup>19</sup>

#### **Peutz-Jeghers Syndrome**

PJS is an autosomal dominant condition mainly characterized by multiple hamartomatous and adenomatous gastrointestinal polyps, predominantly



# NCCN Guidelines Version 2.2021

## Small Bowel Adenocarcinoma

located in the jejunum and ileum.<sup>19,20</sup> A majority of PJS cases occur due to mutations in the *STK11 (LKB1)* gene.<sup>21,22</sup> However, other genetic mutations may be involved, as an estimated half of patients with PJS do not have detectable *STK11/LKB1* mutations.<sup>23</sup> SBA can arise from either hamartomatous or adenomatous polyps, predisposing patients with PJS to SBA. The relative risk of developing SBA has been estimated as 520 compared to the general population,<sup>24</sup> and the lifetime risk of SBA has been estimated between 1.7% to 13% for individuals with PJS.<sup>19,24,25</sup>

### Lynch Syndrome

Lynch syndrome is a hereditary syndrome resulting from germline mutations in DNA mismatch repair (MMR) genes (*MLH1*, *MSH2*, *MSH6*, and *PMS2*). Individuals with Lynch syndrome are estimated to have a lifetime risk of 4% of developing SBA, representing a relative risk of more than 100 compared to the general population.<sup>26,27</sup> Although identifying a mutation in an MMR gene through germline sequencing is definitive for Lynch syndrome, patients usually undergo selection for screening by considering family history and performing an initial test on tumor tissue before sequencing. One of two (or both) different initial tests can be performed on SBA specimens to identify individuals who might have Lynch syndrome: 1) immunohistochemical analysis for MMR protein expression, which is often diminished because of mutation; or 2) polymerase chain reaction (PCR) analysis for microsatellite instability (MSI), which results from MMR deficiency and is detected as changes in the length of repetitive DNA elements in tumor tissue caused by the insertion or deletion of repeated units.<sup>28</sup>

Many NCCN Member Institutions and other comprehensive cancer centers now perform immunohistochemistry (IHC) and sometimes MSI testing on all newly diagnosed CRC and endometrial cancers regardless of family history to determine which patients should have genetic testing for Lynch syndrome.<sup>29-32</sup> This approach may also be applied to patients with a

personal history of SBA, particularly since it has been reported that SBA has a higher percentage of MSI-high (MSI-H)/ MMR-deficient (dMMR) tumors compared to CRC.<sup>33,34</sup> The NCCN Colon/Rectal/Anal Cancers Panel endorses universal MMR or MSI testing of all patients with a personal history of SBA to identify individuals with Lynch syndrome. This testing is also relevant for treatment selection in stage IV disease (see *Pembrolizumab or Nivolumab ± Ipilimumab [for dMMR/MSI-H tumors] as Subsequent-line Therapy*, below). A more detailed discussion is available in the [NCCN Guidelines for Colorectal Cancer Screening](#).

### Clinical Presentation and Workup

The treatment recommendations in this guideline only refer to SBA. For GIST, see the [NCCN Guidelines for Soft Tissue Sarcoma](#); for neuroendocrine tumors, see the [NCCN Guidelines for Neuroendocrine and Adrenal Tumors](#); and for small bowel lymphomas see the [NCCN Guidelines for B-Cell Lymphomas](#).

Most cases of SBA arise in the duodenum, accounting for ~52% to 57% of cases. The remainder arise in the jejunum (18%–29%), ileum (10%–13%), or in an unspecified location of the small bowel (4%–14%).<sup>35-37</sup> Patients with SBA tend to be younger at diagnosis and often present with a higher stage and grade compared to those with CRC.<sup>38</sup> SBA often presents with a local complication of the tumor, most often gastric outlet obstruction in the case of a duodenal SBA or cramping abdominal pain in the case of a jejunal or ileal SBA.<sup>35,39,40</sup> Occult gastrointestinal bleeding is another common presentation for SBA, occurring in approximately one-quarter to one-third of cases.

Patients who present with small bowel cancer require a complete staging workup, including biopsy (if appropriate), pathologic tissue review, imaging studies (see *Small Bowel Imaging*, below), complete blood count (CBC), chemistry profile, carbohydrate antigen 19-9 (CA 19-9), and



# NCCN Guidelines Version 2.2021

## Small Bowel Adenocarcinoma

carcinoembryonic antigen (CEA). Depending on the tumor's location and the patient's history, studies for celiac disease or IBD may be indicated. As discussed above, MMR or MSI testing is recommended for all patients with SBA as MMR/MSI status can function as a prognostic and/or predictive marker and can help identify patients who should be tested for Lynch syndrome (see *Risk Factors for Small Bowel Adenocarcinoma*, above).

### Imaging and Endoscopy

Esophagogastroduodenoscopy (EGD) with endoscopic ultrasound (EUS) is recommended during initial workup and staging for detection and pathologic sampling when a duodenal malignancy is suspected. If obstruction is detected during imaging, palliative diversion or stenting may be considered.<sup>41</sup> EUS is useful for pre-therapeutic staging of proximal small bowel malignancies and may be used to discern duodenal lesions from ampullary, biliary, or pancreatic primaries.<sup>42</sup>

Other endoscopic techniques that are not required for routine staging, but may be useful in certain circumstances, include double balloon endoscopy and capsule endoscopy. A number of studies, both prospective and retrospective, have reported on the effectiveness and safety of double balloon endoscopy for workup of patients with small bowel cancer.<sup>43-45</sup> Specifically, the use of this method may be of particular benefit for patients with small bowel strictures.<sup>46</sup> While capsule endoscopy allows for a more detailed examination of the entire small bowel mucosa, possibly resulting in the diagnosis of SBA when other imaging methods have failed to reveal a primary lesion, it is not the preferred method for initial workup due to its inability to biopsy tissue for diagnosis.<sup>47-49</sup> In the case of a small bowel obstruction or stricture, the capsule may not be excreted naturally, requiring surgical removal. Therefore, capsule endoscopy is contraindicated for these conditions.<sup>41</sup>

CT or magnetic resonance (MR) imaging may be used during initial workup of SBA to evaluate the extent of local tumor invasion and to assess for distant metastases. CT or MR enterography or enteroclysis, techniques involving administration of enteric contrast agent to the gastrointestinal system via oral intake or nasogastric tube, respectively, may improve imaging of the small bowel and, therefore, may be considered when conventional CT or MR with contrast have failed to discern a tumor.<sup>50-53</sup> A prospective study comparing CT enterography to MR enterography in 150 patients with suspected small bowel disease, but negative findings on endoscopy, reported that MR enterography was more accurate than CT enterography, particularly for neoplastic diseases ( $P = .0412$ ).<sup>54</sup>

While PET/CT has not been formally evaluated for ability to detect metastatic SBA, or compared to MR or CT, there have been reports describing its usefulness for this disease.<sup>55</sup> Therefore, while PET/CT is not routinely indicated, it may be considered when CT or MR results are equivocal.

See *Posttreatment Surveillance*, below, for the use of these imaging methods for posttreatment surveillance and for use in individuals with inflammatory bowel disease, celiac disease, or familial syndromes.

### Pathology and Staging

SBA staging is based on the TNM staging system.<sup>6</sup> In the 8<sup>th</sup> edition of the AJCC Staging Manual, T1 tumors involve the lamina propria or submucosa; T2 tumors penetrate through the submucosa into the muscularis propria; T3 tumors penetrate through the muscularis propria into the subserosa or extend into nonperitonealized perimuscular tissue; and T4 tumors perforate the visceral peritoneum or directly invade other organs or structures. Regional lymph node classification includes N0 (no regional lymph node metastasis), N1 (1–2 positive lymph nodes), and N2



# NCCN Guidelines Version 2.2021

## Small Bowel Adenocarcinoma

(3 or more positive nodes). SBA is classified as M1 when distant metastasis is present.<sup>6</sup>

SBA is staged as I or II when a tumor is present without regional lymph node or distant metastases (any T, N0, M0). Stage III disease includes disease with regional lymph node, but not distant, metastasis (any T, N1-2, M0). Stage IV is distant metastatic disease (any T, any N, M1).<sup>6</sup> A number of sources have reported stage III or IV SBA as having significantly worse outcomes compared to earlier stage disease.<sup>6,56,57</sup>

Other factors that may be useful for prognostication, but not used for staging, include the primary tumor site (ie, duodenum, jejunum, ileum); histologic grade; number of lymph nodes evaluated; margin status; lymphovascular invasion; MSI/MMR status; evidence/presence of celiac or IBD; and presence of polyps.<sup>6</sup> The NCCN Panel recommends reporting of these parameters during pathologic review.

### Lymph Node Evaluation

Regional lymph nodes differ based on the site of the primary tumors: retropancreatic, hepatic artery, inferior pancreaticoduodenal, and superior mesenteric nodes are regional to the duodenum; cecal or ileocolic (terminal ileum only, superior mesenteric, or mesenteric [not otherwise specified]) nodes are regional to the jejunum and ileum.

Multiple analyses of patients with SBA in the SEER database have found that longer survival following resection is strongly associated with a lower ratio of positive-to-negative lymph nodes as well as with a higher number of regional lymph nodes assessed during surgery.<sup>38,58-60</sup> Two of these analyses, which considered duodenal and jejunoileal adenocarcinomas separately, concluded that, for adequate staging, a minimum of 5 lymph nodes should be retrieved for duodenal tumors and a minimum of 9 lymph nodes for jejunal or ileal tumors.<sup>59,60</sup> Analyses that pooled duodenal and jejunoileal tumors found that 8 regional lymph nodes should be assessed

for adequate staging,<sup>38,58</sup> although some data have suggested that harvesting even higher numbers of lymph nodes may better predict SBA survival outcomes.<sup>61</sup> Based on these studies, NCCN recommends that a goal for all SBA resections should be the retrieval of at least 8 regional lymph nodes for evaluation.

### Treatment of Stage I-III Small Bowel Adenocarcinoma

#### Surgical Management of Localized Resectable Disease

For local (stage I-III) SBA, primary treatment consists of surgical resection with en bloc removal of the regional lymph nodes. While there have been no prospective, randomized trials to inform surgical technique, retrospective reviews on the subject have been published.<sup>62-64</sup> Intraoperative staging of the abdomen—particularly including the mesentery, omentum, and peritoneum—should be completed in all cases.

The type of resection used to treat localized SBA depends on the location of the primary tumor. Segmental resection of the small bowel is often the mainstay of treatment, although duodenal tumors may require either pancreaticoduodenectomy or segmental duodenal resection. For tumors of the jejunum or ileum, segmentectomy is the preferred method of resection.

Pancreaticoduodenectomy, also known as the Whipple procedure, should be considered for all duodenal cancers, and is particularly appropriate for those arising in the second portion of the duodenum or invading into any portion of the ampulla or pancreas. Minimally invasive procedures, such as laparoscopic surgery, may be considered for pancreaticoduodenectomy, but should only be employed by experienced surgeons. Limited segmentectomy may be considered in SBA involving the third and fourth segments of the duodenum and on the anti-mesenteric side of the intestine, although this approach is controversial based on reports of lower lymph node yields.<sup>64</sup> However, a retrospective study of 1,611 patients with duodenal adenocarcinoma found that patients who



## NCCN Guidelines Version 2.2021 Small Bowel Adenocarcinoma

were treated with radical resection did not show an improvement in overall survival (OS) or disease-specific survival compared to a simple removal of the primary site, after controlling for confounding factors.<sup>65</sup> This finding was despite greater lymph node retrieval with radical resection. Case reports have suggested that segmentectomy and other limited resection methods may be considered for lesions in the first portion of the duodenum, particularly for those <2 cm in size and located on the mesenteric side of the intestine.<sup>63</sup>

NCCN recommends that a goal for all SBA resections should be the retrieval of at least 8 regional lymph nodes for evaluation based on the strong prognostic impact of lymph node metastases and studies showing improved outcomes with higher numbers of lymph nodes assessed during surgery.<sup>58-60</sup> See *Lymph Node Evaluation*, above, for more information on pathologic review of dissected lymph nodes.

### Adjuvant Therapy

Localized SBAs are treated with surgical resection, but local and distant recurrences are common and optimal perioperative therapy is unknown.<sup>66</sup> Therefore, participation in a clinical trial is preferred for all patients with SBA who are considering adjuvant therapy. For discussion of neoadjuvant therapy, see *Primary Treatment of Unresectable Disease*, below.

The ongoing, international phase III BALLAD trial is the first prospective trial investigating the role of adjuvant 5-FU/leucovorin (5-FU/LV) or 5-FU/LV plus oxaliplatin (FOLFOX) compared to observation alone for patients with stage I-III SBA.<sup>67,68</sup> Until the results of BALLAD have been reported, the potential benefits of adjuvant therapy for SBA can be estimated only through retrospective reports. The data from retrospective studies or meta-analyses that have sought to assess the efficacy of adjuvant therapy (either chemotherapy or chemoradiotherapy) for SBA have been mixed, with some showing a benefit to adjuvant therapy,<sup>69-71</sup>

some showing no benefit,<sup>36,72,73</sup> and some showing an equivocal or non-significant benefit.<sup>74,75</sup> Data supporting the use of adjuvant chemoradiation are especially limited, with a recent retrospective review of patients with resected, nonmetastatic duodenal adenocarcinoma showing that patients who received adjuvant chemoradiotherapy (n = 550) had no significant improvement in survival compared to those who received chemotherapy alone (n = 694), even in high-risk cases.<sup>76</sup> Therefore, chemoradiation should be considered only in highly selected patients.

### MSI/MMR Status for Adjuvant Therapy

MSI/MMR is an important piece of information to consider when deciding whether to use adjuvant chemotherapy in patients with stage II SBA. Mutation of MMR genes or modifications of these genes (eg, methylation) can result in MMR protein deficiency and MSI.<sup>77</sup> Tumors showing the presence of MSI are classified as either MSI-H or MSI-low (MSI-L), depending on the extent of instability in the markers tested, whereas tumors without this characteristic are classified as microsatellite-stable (MSS).<sup>78</sup> Patients determined to have dMMR status are biologically the same population as those with MSI-H status.

Data from several large studies in colon cancer have shown that MSI-H (ie, dMMR) tumors have a decreased likelihood to metastasize and that MSI-H/dMMR may function as a prognostic marker for favorable outcomes in stage II disease.<sup>79-81</sup> Some of these same studies also show that a dMMR/MSI-H tumor status may be a predictive marker of decreased benefit and possibly a detrimental impact from adjuvant therapy in patients with stage II colon cancer.<sup>80-82</sup> However, a recent study of 1913 patients with stage II CRC from the QUASAR study, half of whom received adjuvant chemotherapy, showed that although dMMR was prognostic, it did not predict benefit or detrimental impact of chemotherapy.<sup>83</sup> A study of patients in the CALGB 9581 and 89803 trials came to a similar conclusion, though notably this used older, non-standard chemotherapy regimens.<sup>84</sup>





# NCCN Guidelines Version 2.2021

## Small Bowel Adenocarcinoma

Extrapolating from these colon cancer data, patients with stage II MSI-H/dMMR SBA may have a good prognosis and the benefit from adjuvant therapy is unclear.

### ***NCCN Recommendations for Adjuvant Therapy***

Based on the limited data available from retrospective studies of SBA, and extrapolation from studies of colon cancer the NCCN Panel recommends:

- Six months of adjuvant treatment with FOLFOX, capecitabine plus oxaliplatin (CAPEOX), 5-FU/LV, or capecitabine for any locally advanced SBA with positive lymph nodes (stage III). Chemoradiation with capecitabine or infusional 5-FU is another option for stage III duodenal cancer that is margin-positive following resection.
- Observation or 6 months of adjuvant treatment with FOLFOX, CAPEOX, 5-FU/LV, or capecitabine for stage II tumors that are MSS or MMR proficient (pMMR) and have high-risk features. High-risk features include T4 stage, close or positive surgical margins, few lymph nodes examined (<5 for duodenal or <8 for jejunal/ileal primary tumor location), or tumor perforation.<sup>38,58,72</sup> Studies in CRC, and a retrospective report in SBA, have identified lymphovascular or perineural invasion and poorly differentiated histology as poor prognostic factors.<sup>72,85-87</sup> Therefore, adjuvant therapy may be considered for patients with these factors as well. Chemoradiation with capecitabine or infusional 5-FU is another option for duodenal cancer that meets these criteria and is margin-positive following resection.
- Observation or 6 months of adjuvant treatment with 5-FU/LV or capecitabine for T3, N0, M0 (stage IIA) tumors that are MSS or pMMR and have no high-risk features.

- Observation following surgical treatment for all stage I tumors and for stage II tumors that are MSI-H or dMMR.

Due to poorer survival of stage III SBA compared to CRC, and the fact that the trial enrolled no patients with SBA, extrapolation from the IDEA collaboration (wherein 3 months of fluoropyrimidine/oxaliplatin was shown to be non-inferior to 6 months of therapy for CRC)<sup>88</sup> is not currently recommended for SBA.

### **Primary Treatment of Unresectable Disease**

For some patients with locally unresectable or medically inoperable SBA, conversion to resectable disease may be a goal. A limited amount of data has demonstrated that neoadjuvant therapy may be beneficial in converting unresectable SBA to resectable disease. A retrospective study of patients with unresectable or recurrent duodenal adenocarcinoma who were treated with neoadjuvant chemotherapy or chemoradiation found that 9 out of 10 patients showed conversion to resectable disease following neoadjuvant therapy. At the time of data collection, 5 patients were still alive (ranging from 18–83 months postoperatively), suggesting prolonged survival following conversion to resectable disease.<sup>89</sup> In addition, neoadjuvant chemoradiation was studied in two small prospective trials. A phase II trial including patients with duodenal or pancreatic adenocarcinomas reported that 4 of 5 patients with tumors in the duodenum were able to undergo resection following neoadjuvant chemoradiation.<sup>90</sup> Another small prospective study of patients with duodenal or pancreatic adenocarcinomas reported that all 4 patients with duodenal cancer underwent curative resection following neoadjuvant chemoradiation and experienced a complete pathologic response.<sup>91</sup>

Since many small bowel cancers present at an advanced stage, malignant small bowel obstruction is a common complication. One retrospective Eastern European study reported that most patients with small bowel



# NCCN Guidelines Version 2.2021

## Small Bowel Adenocarcinoma

cancer presented due to an emergency situation,<sup>39</sup> with obstruction being a common complication for SBA, accounting for 22% to 57.9% of these cases.<sup>39,92-94</sup> Malignant small bowel obstruction may be treated palliatively with either surgical diversion or stenting. While most of the literature on palliative treatment of malignant small bowel obstruction comes from pancreatic cancer, there are a few studies that include SBA cases.<sup>39,95-97</sup> One retrospective study concluded that there was no difference in post-stent survival between patients with pancreatic and nonpancreatic cancers, and that patients with nonpancreatic cancers (including SBA) showed a longer OS.<sup>95</sup>

Based on these data, the panel recommends that patients with locally unresectable or medically inoperable SBA may undergo neoadjuvant therapy, during which they should be routinely monitored for conversion to resectable disease. Neoadjuvant chemoradiation may be indicated for duodenal disease that remains unresectable following a course of induction chemotherapy, but is controversial and should be considered on an individual case basis. Alternatively, in cases where conversion to resectable disease is not feasible, palliative chemotherapy may be considered. Palliative diversion or stenting is recommended if a small bowel obstruction is present.

### Treatment of Distant Metastatic (Stage IV) SBA

Approximately 32% of patients diagnosed with SBA have stage IV (distant metastatic) disease.<sup>38</sup> The most common sites for metastatic spread include the peritoneal cavity and liver, consistent with other gastrointestinal malignancies.<sup>6</sup> While 5-year survival is relatively high (85%) for localized disease, patients with stage IV SBA have a 5-year relative survival of only 42%.<sup>98</sup> In addition, recurrence rates of localized SBA treated with surgery are high, with many of these patients developing distant metastases.<sup>35</sup> The NCCN recommendations for treatment of stage IV SBA are discussed below.

### Metastasectomy

While resectable metastases are rare for SBA and the data supporting metastasectomy for SBA are limited, a retrospective analysis of patients with non-CRC, nonendocrine liver metastases (including 28 patients with small bowel cancers and 12 patients with duodenal cancers) showed promising survival rates following resection of liver metastases.<sup>99</sup> The 5-year survival rate for small bowel cancers was 49% with a median survival of 58 months. For duodenal cancers, the 5-year survival rate was 21% with a median survival of 34 months. Recently, another retrospective study of 34 patients undergoing resection of SBA metastases reported a median OS of 28.2 months and a relapse-free survival of 18.7 months.<sup>100</sup> 41.2% of patients in this study survived more than 3 years. 88.2% of patients in this study received perioperative chemotherapy and poor differentiation, invaded margins, and lymphatic invasion of the primary tumor were identified as poor prognostic factors. Therefore, certain patients with SBA and limited metastasis to visceral organs may be candidates for metastasectomy. If metastasectomy is being considered, a multidisciplinary team, including a surgeon experienced in the resection of metastases, should be consulted.

### Peritoneal Carcinomatosis

Peritoneal carcinomatosis (peritoneal metastases) has been shown to affect 25% to 50% of patients with stage IV SBA. Peritoneal carcinomatosis occurs more frequently in tumors arising from the jejunum or ileum and less commonly in duodenal tumors.<sup>101</sup> Peritoneal carcinomatosis generally carries a poor prognosis with a reported median OS of 5.9 months.<sup>102</sup> The goal of treatment for unresectable peritoneal metastases is palliative and primarily consists of systemic therapy (see *Systemic Therapy for Metastatic Disease*).

For resectable peritoneal carcinomatosis, surgical cytoreduction may be considered. For peritoneal metastases that present synchronously with the



# NCCN Guidelines Version 2.2021

## Small Bowel Adenocarcinoma

primary tumor, resection of the primary and cytoreduction of peritoneal metastases may be carried out concurrently. A multidisciplinary team evaluation at an experienced center is important if considering this treatment approach. Data supporting the use of hyperthermic intraperitoneal chemotherapy (HIPEC) for SBA with peritoneal carcinomatosis are extremely limited, consisting entirely of small, retrospective studies.<sup>102-107</sup> In addition, the recent phase III PRODIGE 7 study showed no benefit of oxaliplatin-based HIPEC in CRC patients compared to cytoreduction alone.<sup>108</sup> Significant morbidity and mortality are associated with this procedure. Various studies have reported morbidity rates ranging from 19% to 31% for serious adverse events and mortality rates ranging from 0% to 4%.<sup>103-107</sup> Furthermore, recurrences after the procedure are common.<sup>106,107</sup> Based on this lack of evidence, HIPEC cannot be recommended for this population unless more robust data become available.

### Systemic Therapy for Metastatic Disease

Data supporting systemic therapy for advanced adenocarcinoma of the small bowel were also almost entirely limited to retrospective reports,<sup>109-112</sup> although recently several small phase II trials for SBA have been reported. Based on the results from these studies, several systemic therapy regimens are recommended for treatment of metastatic SBA. However, participation in clinical trials is especially encouraged for patients with SBA based on the lack of data.

The choice of therapy is based on consideration of the goals of therapy, the type and timing of prior therapy, and the differing toxicity profiles of the constituent drugs. Furthermore, an evaluation of the efficacy and safety of these regimens for an individual patient must take into account the performance status of the patient. As initial therapy for advanced disease in a patient appropriate for intensive therapy (ie, one with a good tolerance for this therapy for whom a high tumor response rate would be potentially

beneficial) without prior platinum resistance, the panel recommends a choice of 3 chemotherapy regimens: FOLFOX, CAPEOX, or FOLFOXIRI (infusional 5-FU, LV, oxaliplatin, irinotecan); any of which may be combined with bevacizumab. For patients who are not appropriate for intensive therapy, treatment options would exclude the more toxic components of these regimens with 5-FU/LV or capecitabine with or without bevacizumab recommended as first-line therapy for these patients.

The choice of second-line therapy depends on the MMR/MSI status of the tumor. For tumors that are dMMR or MSI-H, checkpoint inhibitor therapy with anti-PD-1 inhibitors, alone or in combination with an anti-CTLA4 inhibitor, is recommended in the second-line setting. FOLFIRI or taxane-based chemotherapies are options in the second line for pMMR/MSS tumors, or those that are refractory to checkpoint inhibitor therapies. Larotrectinib is an option in subsequent lines of therapy for metastatic SBA with neurotrophic tyrosine receptor kinase (*NTRK*) gene fusion and no satisfactory alternative treatments.

### Genetic Alterations in SBA

Emerging research has shown that SBA has a distinct genetic profile, which sets it apart from CRC or gastroesophageal cancers, the two cancer types SBA is most often likened to. While *KRAS* and *TP53* alterations are frequently identified in both SBA and CRC, *APC* mutations are significantly less common in SBA (27% in SBA vs. 76% in CRC;  $P < .001$ ).<sup>34</sup> Considering the near ubiquity of *APC* mutation and its well-established role in CRC carcinogenesis, this suggests that neoplastic transformation in SBA is unique compared to CRC.<sup>33,34</sup>

*SMAD4* and *CDKN2A* mutations are more commonly seen compared to gastroesophageal cancers and CRC. Though *BRAF* mutations occur at a similar rate as seen in CRC, only 10% of *BRAF*-mutant SBAs have a V600E alteration, compared with >70% in *BRAF*-mutant CRC.<sup>34</sup> Importantly, human epidermal growth factor receptor 2 (HER2) alterations,



# NCCN Guidelines Version 2.2021

## Small Bowel Adenocarcinoma

MSI-H/dMMR, programmed death-ligand 1 (PD-L1) expression, and high tumor mutational burden are enhanced in SBA compared to CRC,<sup>34,113-115</sup> and may reveal greater importance of targeted or immunotherapeutic treatments compared to current CRC treatment algorithms.

### **Regimens Not Recommended for SBA**

While many of the systemic therapy regimens recommended for treatment of metastatic SBA are extrapolated from data for CRC, there are several regimens commonly used for metastatic CRC that are not recommended for SBA based either on a lack of data supporting their use or data suggesting that these regimens do not work for metastatic SBA.

A 2017 retrospective analysis reported that the efficacy of cetuximab-containing chemotherapy for *RAS* wild-type SBA was inconclusive.<sup>116</sup> Subsequently, a phase II trial published in 2018 showed that panitumumab has no clinically meaningful activity in *RAS* wild-type SBA;<sup>117</sup> therefore, cetuximab or panitumumab should not be used for treatment of SBA.

While trifluridine-tipiracil or regorafenib are recommended as subsequent therapy options for metastatic CRC, there are no data to support their use for SBA and are, therefore, not recommended.

### **FOLFOX or CAPEOX as First-line Therapy**

Both FOLFOX and CAPEOX have been evaluated prospectively for first-line treatment of advanced SBA in phase II clinical trials. One of these trials evaluated CAPEOX in 30 patients with advanced adenocarcinomas of the small bowel and ampulla of Vater. The overall response rate (ORR) (the primary endpoint) was 50%, with 10% achieving complete response.<sup>118</sup> A similar response rate of 48.5% (95% CI, 31%–67%) was seen in another small phase II study of 33 patients that assessed the efficacy of FOLFOX in first-line treatment of advanced SBA.<sup>119</sup> Likewise, another phase II study reported an ORR of 45% for 24 patients with metastatic or unresectable SBA who were treated with FOLFOX, with a

median progression-free survival (PFS) and OS of 5.9 and 17.3 months, respectively.<sup>120</sup> These response rates to CAPEOX and FOLFOX were much higher than the 18% response rate seen in another small phase II study that evaluated 5-FU/doxorubicin/mitomycin C in patients with metastatic SBA.<sup>121</sup> Adverse events reported across these three trials were similar, with neutropenia, thrombocytopenia, nausea, vomiting, diarrhea, peripheral neuropathy, and fatigue reported most frequently.<sup>118-120</sup> Retrospective studies have supported the results of these trials, reporting that the combination of a fluoropyrimidine with oxaliplatin was the most effective first-line therapy for advanced SBA.<sup>111,122,123</sup> Based on these data, FOLFOX or CAPEOX are recommended as first-line therapy options for treatment of patients with advanced SBA who are appropriate for intensive therapy.

### **FOLFOXIRI as First-line Therapy**

While the role of FOLFOXIRI for treatment of SBA has not been formally evaluated, CAPIRINOX (capecitabine, irinotecan, oxaliplatin) has been tested as first-line treatment in a phase II trial of 33 patients with advanced SBA.<sup>124</sup> In this trial, CAPIRINOX—dose-adjusted according to *UGT1A1* genotype—showed a response rate of 37.5% (95% CI, 21%–56%), with a median PFS and OS of 8.9 and 13.4 months, respectively. Neither hematologic toxicity nor tumor response rate differed significantly by *UGT1A1* genotype, supporting the feasibility of genotype-directed dosing for CAPIRINOX. The NCCN Panel does not recommend use of CAPIRINOX for SBA due to concerns about toxicity, but the recommendation for FOLFOXIRI is extrapolated from the results of this study.

### **FOLFOX, CAPEOX, or FOLFOXIRI Plus Bevacizumab as First-line Therapy**

While data supporting the addition of biologics to FOLFOX, CAPEOX, or FOLFOXIRI are currently extremely limited, a single-phase II trial has reported that CAPEOX in combination with bevacizumab is safe and



# NCCN Guidelines Version 2.2021

## Small Bowel Adenocarcinoma

efficacious in patients with SBA.<sup>125</sup> Retrospective analyses have supported these results, reporting favorable outcomes in patients treated with bevacizumab-containing chemotherapy regimens without adding significant toxicity.<sup>116,126</sup> Based on these data, FOLFOX, CAPEOX, or FOLFOXIRI may be given with or without bevacizumab as first-line therapy for advanced SBA.

### ***Pembrolizumab or Nivolumab ± Ipilimumab (for dMMR/MSI-H tumors) as Subsequent-line Therapy***

Pembrolizumab is a PD-1 inhibitor that was evaluated as a subsequent-line therapy for treatment-refractory metastatic cancers in a phase 2 study that included 3 cohorts: 1) dMMR colorectal adenocarcinomas, 2) MMR-proficient colorectal adenocarcinomas, and 3) dMMR cancers of types other than CRC.<sup>127</sup> This third cohort included 2 patients with small bowel cancers. The immune-related objective response rate and immune-related PFS rate were 40% and 78%, respectively, for patients with dMMR CRC and 71% and 67% for patients with dMMR non-CRC. Common adverse events of clinical interest included rash or pruritus; thyroiditis, hypothyroidism, or hypophysitis; and asymptomatic pancreatitis.<sup>127</sup> Based on the results of this study, the FDA granted accelerated approval to pembrolizumab in May 2017 for patients with unresectable or metastatic dMMR or MSI-H solid tumors that have progressed following prior treatment and have no satisfactory alternative treatment options.<sup>128</sup> More recently, an abstract reported results of ZEBRA, a multicenter, phase 2 study of pembrolizumab in patients with previously treated, advanced SBA.<sup>129</sup> The results of this study confirmed efficacy of pembrolizumab for dMMR/MSI-H SBA. Furthermore, while pembrolizumab did not achieve the goal ORR for this study, there was some evidence that this therapy may control disease in some patients with MSS SBA. Of 18 patients with confirmed MSS SBA, there was a 50% disease control rate, although further study is needed to confirm this result.<sup>129</sup>

Another PD-1 inhibitor, nivolumab—alone or in combination with the CTLA-4 inhibitor, ipilimumab—has been studied in patients with dMMR metastatic CRC in the phase II, multi-cohort CheckMate-142 trial.<sup>130,131</sup> One cohort of this trial included 74 patients with dMMR CRC who were treated with nivolumab. ORR for these patients was 31.1% (95% CI, 20.8–42.9), with 69% of patients having disease control for at least 12 weeks. Median duration of response had not yet been reached at the time of data collection. PFS and OS were 50% and 73%, respectively, at 1 year. Grade 3 or 4 drug-related adverse events occurred in 20% of patients, with increased amylase and increased lipase being the most common.<sup>130</sup> Another cohort of the CheckMate-142 trial included 119 patients with dMMR CRC who were treated with nivolumab in combination with ipilimumab. For this cohort, ORR was 55% (95% CI, 45.2–63.8) and the disease control rate for at least 12 weeks was 80%. PFS and OS were 71% and 85%, respectively, at 1 year. In addition, significant, clinically meaningful improvements were observed in patient-reported outcomes of functioning, symptoms, and quality of life. Grade 3 to 4 treatment-related adverse events occurred in 32% of patients, but were manageable.<sup>131</sup>

Based on these positive results for CRC, and the data showing benefit of pembrolizumab in SBA, the NCCN Panel recommends either pembrolizumab or nivolumab, with or without ipilimumab, as second-line treatment options for dMMR/MSI-H advanced SBA. SBA has been reported to have a higher incidence of dMMR/MSI-H and higher rates of PD-L1 IHC positivity compared to CRC,<sup>33,34,113</sup> making checkpoint inhibition an important treatment option for some SBA patients.

### ***Taxane-based Chemotherapy as Subsequent-line Therapy***

While almost all of the phase II trials of systemic therapy for SBA have focused on first-line therapy, a phase II trial including 13 patients with SBA studied the efficacy of nab-paclitaxel in the refractory disease setting.<sup>132</sup> Patients with SBA in this trial had received a median of 2 prior lines of



# NCCN Guidelines Version 2.2021

## Small Bowel Adenocarcinoma

therapy including a fluoropyrimidine and oxaliplatin. Of the 10 patients with SBA who were evaluable for efficacy, 2 showed a partial response to nab-paclitaxel and an additional 3 had stable disease per RECIST criteria, yielding a disease control rate of 50%. Common grade 3 or 4 toxicities across the entire study population included fatigue (12%), neutropenia (9%), febrile neutropenia (9%), dehydration (6%), and thrombocytopenia (6%).<sup>132</sup>

A single-center, retrospective review reported on 20 patients with advanced SBA who were treated with taxane-based therapy (either as single therapy or in combination).<sup>133</sup> Of these cases, 30% showed disease response, 35% showed stable disease, and 35% showed progression. Median time to progression was 3.8 months (95% CI, 2.9–4.6) and median OS was 10.7 months (95% CI, 3.1–18.3). Based on these data, taxane-based chemotherapy is a recommended option for second- or subsequent-line therapy, although only nab-paclitaxel has prospective, published data to support its use for treatment of SBA.

### ***FOLFIRI as Subsequent-line Therapy***

A retrospective, multicenter study evaluated the efficacy of FOLFIRI as second-line therapy for patients with advanced SBA who had received platinum-based chemotherapy in the first-line setting.<sup>134</sup> Of the 28 patients who fit this treatment paradigm, the ORR was 20% and disease control rate was 52%. The median PFS and OS were 3.2 and 10.5 months. Grade 3–4 toxicity was reported in 48% of patients. Based on these data, FOLFIRI is recommended as a treatment option for second- or subsequent-line treatment of advanced SBA.

### ***Larotrectinib as Subsequent-line Therapy***

A pooled analysis of 3 studies (a phase 1 including adults, a phase 1/2 involving children, and a phase 2 involving adolescents and adults) studied the safety and efficacy of larotrectinib in patients with *NTRK* gene fusion-positive tumors, including 4 patients with colon cancer and 1 with

cancer of the appendix.<sup>135</sup> For the whole population, the ORR was 75% (95% CI, 61%–85%) by independent review and 80% (95% CI, 67%–90%) by investigator assessment. Larotrectinib was found to be well-tolerated as the majority (93%) of adverse events were grades 1 or 2 and no treatment-related adverse events of grades 3 or 4 occurred in more than 5% of patients.<sup>135</sup> Based on these data, the FDA approved larotrectinib for metastatic solid tumors with *NTRK* gene fusion and no satisfactory alternative treatments on November 26, 2018.<sup>136</sup>

### **Posttreatment Surveillance**

After curative-intent surgery and adjuvant chemotherapy, if administered, post-treatment surveillance of patients with SBA is performed to evaluate for possible therapeutic complications, identify disease recurrence, and discover new metachronous neoplasms at a preinvasive stage. A retrospective study of 146 patients with SBA who underwent cancer-directed surgery found that 39% subsequently developed disease recurrence, with a median time to recurrence of 25 months. Of the patients with disease recurrence, 57% developed distant metastases, 19% developed carcinomatosis, 7% recurred in the abdominal wall, and 17% developed local recurrences.<sup>35</sup> Due to the lack of data regarding optimal surveillance following curative-intent treatment of SBA, a similar approach to CRC surveillance is recommended—including history and physical examination; CEA and/or CA 19-9 measurement; and CT of the chest, abdomen, and pelvis. For data supporting the recommended surveillance approach for CRC, see the *Posttreatment Surveillance* section in the [NCCN Guidelines for Colon Cancer](#).

Patients with SBA who were determined to have Crohn's disease or familial syndromes (ie, Lynch, FAP, PJS) may require more intensive surveillance due to their elevated risk of developing further SBAs.<sup>11,12</sup> Endoscopy may be a feasible method for SBA surveillance in patients with Crohn's disease,<sup>12,137</sup> although one prospective study found a low (33%)



# NCCN Guidelines Version 2.2021

## Small Bowel Adenocarcinoma

sensitivity rate for SBA endoscopic screening.<sup>138</sup> A number of studies have supported the use of endoscopy/enteroscopy for small bowel surveillance in patients with Lynch syndrome, FAP, or PJS.<sup>139-146</sup> For further details on endoscopic small bowel evaluation, see the section on *Imaging and Endoscopy* under *Clinical Presentation and Workup*, above.

### Survivorship

Based on the rarity and poor prognosis of SBA, there is a dearth of data regarding survivorship for this disease. The panel recommendations for survivorship are largely extrapolated from the NCCN Guidelines for Colon Cancer, with some specific recommendations included for patients with celiac or Crohn's disease who are at elevated risk of developing additional SBAs.<sup>11,12,14</sup> This section will provide an overview of the panel recommendations for survivorship; for more detailed information, see the *Survivorship* section in the [NCCN Guidelines for Colon Cancer](#).

The panel recommends that a prescription for survivorship and transfer of care to the primary care physician be written.<sup>147</sup> The oncologist and primary care provider should have defined roles in the surveillance period, with roles communicated to the patient. Other recommendations include monitoring for late or long-term sequelae of treatment, such as oxaliplatin-induced peripheral neuropathy, fatigue, pain, sexual dysfunction, and emotional or social distress.<sup>148-152</sup> Specific management interventions to address these and other side effects are described in a review.<sup>153</sup> Disease preventive measures, such as immunizations; early disease detection through periodic screening for second primary cancers (eg, breast, cervical, or prostate cancers); and routine good medical care and monitoring are recommended. The [NCCN Guidelines for Survivorship](#) provide screening, evaluation, and treatment recommendations for common consequences of cancer and cancer treatment to aid health care professionals who work with survivors of adult-onset cancer in the post-

treatment period, including those in specialty cancer survivor clinics and primary care practices.

### Summary

SBA is a rare malignancy, with a rising incidence in recent decades. Compared to CRC, SBA is more often diagnosed at advanced stages, suggesting the difficulty of detecting these cancers and highlighting the lack of screening programs, even for high-risk individuals. The majority of SBAs arise in the duodenum and are associated with poorer prognosis, with up to a third of resectable patients experiencing early relapse. To date, the only curative therapy for SBA is surgery.

For local disease, segmental resection of the small bowel is the mainstay of treatment, though duodenal tumors may require either pancreaticoduodenectomy or segmental duodenal resection. Database analyses have reported significantly improved outcomes when 8 or more lymph nodes are resected. In addition, the use of radiation therapy for retroperitoneal-based duodenal adenocarcinomas is a complex decision-making process. Fluoropyrimidine-based adjuvant therapy may be considered for some patients with SBA, though no studies have yet shown a definitive benefit of this approach. Results from the international, phase III adjuvant clinical study (BALLAD) investigating observation versus 5-FU versus FOLFOX for patients with resected stage I-III SBA should shed light on this approach in the coming years.

Metastatic SBA may rarely be treated with curative intent via primary tumor resection and metastasectomy; however, most patients with metastatic SBA are treated with systemic therapy. Systemic therapy options include fluoropyrimidine-based chemotherapy, taxane-based chemotherapy, or checkpoint inhibitors. Recently, SBA's unique genetic profile has been a topic of research, which may lead to new targeted or immunotherapeutic treatment options for SBA.



# NCCN Guidelines Version 2.2021

## Small Bowel Adenocarcinoma

### References

1. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2019. *CA Cancer J Clin* 2019;69:7-34. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30620402>.
2. Noone AM, Howlader N, Krapcho M, et al. SEER Cancer Statistics Review, 1975-2015, based on November 2017 SEER data submission, posted to the SEER web site, April 2018. Bethesda, MD: National Cancer Institute; 2018. Available at: [https://seer.cancer.gov/csr/1975\\_2015/](https://seer.cancer.gov/csr/1975_2015/).
3. Li J, Saif MW. Small Bowel Adenocarcinoma. In: Saif MW, ed. *Gastrointestinal Malignancies*. New York, NY: Demos Medical Publishing, LLC; 2010:171-176.
4. Bilimoria KY, Bentrem DJ, Wayne JD, et al. Small bowel cancer in the United States: changes in epidemiology, treatment, and survival over the last 20 years. *Ann Surg* 2009;249:63-71. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19106677>.
5. Locher C, Batumona B, Afchain P, et al. Small bowel adenocarcinoma: French intergroup clinical practice guidelines for diagnosis, treatments and follow-up (SNFGE, FFCD, GERCOR, UNICANCER, SFCD, SFED, SFRO). *Dig Liver Dis* 2018;50:15-19. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29174568>.
6. Amin MB, Edge SB, Greene F, et al., eds. *AJCC Cancer Staging Manual*, 8th ed. New York: Springer International Publishing; 2017.
7. U.S. National Library of Medicine-Key MEDLINE® Indicators. Available at: [http://www.nlm.nih.gov/bsd/bsd\\_key.html](http://www.nlm.nih.gov/bsd/bsd_key.html). Accessed October 8, 2018.
8. Bennett CM, Coleman HG, Veal PG, et al. Lifestyle factors and small intestine adenocarcinoma risk: A systematic review and meta-analysis. *Cancer Epidemiol* 2015;39:265-273. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25736860>.
9. Boffetta P, Hazelton WD, Chen Y, et al. Body mass, tobacco smoking, alcohol drinking and risk of cancer of the small intestine--a pooled analysis of over 500,000 subjects in the Asia Cohort Consortium. *Ann Oncol* 2012;23:1894-1898. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22147734>.
10. Algaba A, Guerra I, Marin-Jimenez I, et al. Incidence, management, and course of cancer in patients with inflammatory bowel disease. *J Crohns Colitis* 2015;9:326-333. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25687203>.
11. Cahill C, Gordon PH, Petrucci A, Boutros M. Small bowel adenocarcinoma and Crohn's disease: any further ahead than 50 years ago? *World J Gastroenterol* 2014;20:11486-11495. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25206256>.
12. Grolleau C, Pote NM, Guedj NS, et al. Small bowel adenocarcinoma complicating Crohn's disease: a single-centre experience emphasizing the importance of screening for dysplasia. *Virchows Arch* 2017;471:611-617. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28421339>.
13. Laukoetter MG, Mennigen R, Hannig CM, et al. Intestinal cancer risk in Crohn's disease: a meta-analysis. *J Gastrointest Surg* 2011;15:576-583. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21152994>.
14. Rampertab SD, Forde KA, Green PH. Small bowel neoplasia in coeliac disease. *Gut* 2003;52:1211-1214. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12865284>.
15. Benhammane H, El M'rabet F Z, Idrissi Serhouchni K, et al. Small bowel adenocarcinoma complicating coeliac disease: a report of three cases and the literature review. *Case Rep Oncol Med* 2012;2012:935183. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23243535>.
16. Zullo A, De Francesco V, Manta R, et al. A challenging diagnosis of jejunal adenocarcinoma in a celiac patient: Case report and systematic review of the literature. *J Gastrointest Liver Dis* 2017;26:411-415. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29253057>.
17. Galiatsatos P, Foulkes WD. Familial adenomatous polyposis. *Am J Gastroenterol* 2006;101:385-398. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16454848>.





## NCCN Guidelines Version 2.2021 Small Bowel Adenocarcinoma

18. Half E, Bercovich D, Rozen P. Familial adenomatous polyposis. *Orphanet J Rare Dis* 2009;4:22. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19822006>.
19. Shenoy S. Genetic risks and familial associations of small bowel carcinoma. *World J Gastrointest Oncol* 2016;8:509-519. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27326320>.
20. Tomlinson IP, Houlston RS. Peutz-Jeghers syndrome. *J Med Genet* 1997;34:1007-1011. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9429144>.
21. Hemminki A, Markie D, Tomlinson I, et al. A serine/threonine kinase gene defective in Peutz-Jeghers syndrome. *Nature* 1998;391:184-187. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9428765>.
22. Jenne DE, Reimann H, Nezu J, et al. Peutz-Jeghers syndrome is caused by mutations in a novel serine threonine kinase. *Nat Genet* 1998;18:38-43. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9425897>.
23. Lim W, Hearle N, Shah B, et al. Further observations on LKB1/STK11 status and cancer risk in Peutz-Jeghers syndrome. *Br J Cancer* 2003;89:308-313. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12865922>.
24. Giardiello FM, Brensinger JD, Tersmette AC, et al. Very high risk of cancer in familial Peutz-Jeghers syndrome. *Gastroenterology* 2000;119:1447-1453. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11113065>.
25. Hearle N, Schumacher V, Menko FH, et al. Frequency and spectrum of cancers in the Peutz-Jeghers syndrome. *Clin Cancer Res* 2006;12:3209-3215. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16707622>.
26. Koornstra JJ, Kleibeuker JH, Vasen HF. Small-bowel cancer in Lynch syndrome: is it time for surveillance? *Lancet Oncol* 2008;9:901-905. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18760246>.
27. Vasen HF, Wijnen JT, Menko FH, et al. Cancer risk in families with hereditary nonpolyposis colorectal cancer diagnosed by mutation analysis. *Gastroenterology* 1996;110:1020-1027. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8612988>.
28. Hendriks YM, de Jong AE, Morreau H, et al. Diagnostic approach and management of Lynch syndrome (hereditary nonpolyposis colorectal carcinoma): a guide for clinicians. *CA Cancer J Clin* 2006;56:213-225. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16870997>.
29. Beamer LC, Grant ML, Espenschied CR, et al. Reflex immunohistochemistry and microsatellite instability testing of colorectal tumors for Lynch syndrome among US cancer programs and follow-up of abnormal results. *J Clin Oncol* 2012;30:1058-1063. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22355048>.
30. Burt RW. Who should have genetic testing for the Lynch syndrome? *Ann Intern Med* 2011;155:127-128. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21768586>.
31. Matloff J, Lucas A, Polydorides AD, Itzkowitz SH. Molecular tumor testing for Lynch syndrome in patients with colorectal cancer. *J Natl Compr Canc Netw* 2013;11:1380-1385. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24225971>.
32. Ward RL, Hicks S, Hawkins NJ. Population-based molecular screening for Lynch syndrome: implications for personalized medicine. *J Clin Oncol* 2013;31:2554-2562. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23733757>.
33. Aparicio T, Svrcek M, Zaanani A, et al. Small bowel adenocarcinoma phenotyping, a clinicobiological prognostic study. *Br J Cancer* 2013;109:3057-3066. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24196786>.
34. Schrock AB, Devoe CE, McWilliams R, et al. Genomic profiling of small-bowel adenocarcinoma. *JAMA Oncol* 2017;3:1546-1553. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28617917>.



## NCCN Guidelines Version 2.2021 Small Bowel Adenocarcinoma

35. Dabaja BS, Suki D, Pro B, et al. Adenocarcinoma of the small bowel: presentation, prognostic factors, and outcome of 217 patients. *Cancer* 2004;101:518-526. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15274064>.
36. Halfdanarson TR, McWilliams RR, Donohue JH, Quevedo JF. A single-institution experience with 491 cases of small bowel adenocarcinoma. *Am J Surg* 2010;199:797-803. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20609724>.
37. Howe JR, Karnell LH, Menck HR, Scott-Conner C. The American College of Surgeons Commission on Cancer and the American Cancer Society. Adenocarcinoma of the small bowel: review of the National Cancer Data Base, 1985-1995. *Cancer* 1999;86:2693-2706. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10594865>.
38. Overman MJ, Hu CY, Kopetz S, et al. A population-based comparison of adenocarcinoma of the large and small intestine: insights into a rare disease. *Ann Surg Oncol* 2012;19:1439-1445. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22187121>.
39. Negoï I, Paun S, Hostiuc S, et al. Most small bowel cancers are revealed by a complication. *Einstein (Sao Paulo)* 2015;13:500-505. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26676271>.
40. Gustafsson BI, Siddique L, Chan A, et al. Uncommon cancers of the small intestine, appendix and colon: an analysis of SEER 1973-2004, and current diagnosis and therapy. *Int J Oncol* 2008;33:1121-1131. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19020744>.
41. Hara AK, Leighton JA, Sharma VK, et al. Imaging of small bowel disease: comparison of capsule endoscopy, standard endoscopy, barium examination, and CT. *Radiographics* 2005;25:697-711; discussion 711-698. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15888619>.
42. Nylund K, Odegaard S, Hausken T, et al. Sonography of the small intestine. *World J Gastroenterol* 2009;15:1319-1330. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19294761>.
43. Chen WG, Shan GD, Zhang H, et al. Double-balloon enteroscopy in small bowel diseases: Eight years single-center experience in China. *Medicine (Baltimore)* 2016;95:e5104. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27759639>.
44. Cazzato IA, Cammarota G, Nista EC, et al. Diagnostic and therapeutic impact of double-balloon enteroscopy (DBE) in a series of 100 patients with suspected small bowel diseases. *Dig Liver Dis* 2007;39:483-487. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17379586>.
45. Mitsui K, Tanaka S, Yamamoto H, et al. Role of double-balloon endoscopy in the diagnosis of small-bowel tumors: the first Japanese multicenter study. *Gastrointest Endosc* 2009;70:498-504. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19555947>.
46. Sunada K, Yamamoto H, Kita H, et al. Clinical outcomes of enteroscopy using the double-balloon method for strictures of the small intestine. *World J Gastroenterol* 2005;11:1087-1089. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15742422>.
47. Cobrin GM, Pittman RH, Lewis BS. Increased diagnostic yield of small bowel tumors with capsule endoscopy. *Cancer* 2006;107:22-27. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16736516>.
48. Bailey AA, Debinski HS, Appleyard MN, et al. Diagnosis and outcome of small bowel tumors found by capsule endoscopy: a three-center Australian experience. *Am J Gastroenterol* 2006;101:2237-2243. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17032187>.
49. Cheung DY, Lee IS, Chang DK, et al. Capsule endoscopy in small bowel tumors: a multicenter Korean study. *J Gastroenterol Hepatol* 2010;25:1079-1086. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20594222>.
50. Boudiaf M, Jaff A, Soyer P, et al. Small-bowel diseases: prospective evaluation of multi-detector row helical CT enteroclysis in 107 consecutive patients. *Radiology* 2004;233:338-344. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15459329>.



# NCCN Guidelines Version 2.2021

## Small Bowel Adenocarcinoma

51. Soyer P, Aout M, Hoeffel C, et al. Helical CT-enteroclysis in the detection of small-bowel tumours: a meta-analysis. *Eur Radiol* 2013;23:388-399. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22865269>.
52. Cronin CG, Lohan DG, Browne AM, et al. Magnetic resonance enterography in the evaluation of the small bowel. *Semin Roentgenol* 2009;44:237-243. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19715789>.
53. Masselli G, Casciani E, Poletti E, et al. Magnetic resonance imaging of small bowel neoplasms. *Cancer Imaging* 2013;13:92-99. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23524074>.
54. Masselli G, Di Tola M, Casciani E, et al. Diagnosis of small-bowel diseases: Prospective comparison of multi-detector row CT enterography with MR enterography. *Radiology* 2016;279:420-431. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26599801>.
55. Cronin CG, Scott J, Kambadakone A, et al. Utility of positron emission tomography/CT in the evaluation of small bowel pathology. *Br J Radiol* 2012;85:1211-1221. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22919004>.
56. Sakae H, Kanzaki H, Nasu J, et al. The characteristics and outcomes of small bowel adenocarcinoma: a multicentre retrospective observational study. *Br J Cancer* 2017;117:1607-1613. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28982111>.
57. Young JI, Mongoue-Tchokote S, Wieghard N, et al. Treatment and survival of small-bowel adenocarcinoma in the United States: A comparison with colon cancer. *Dis Colon Rectum* 2016;59:306-315. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26953989>.
58. Overman MJ, Hu CY, Wolff RA, Chang GJ. Prognostic value of lymph node evaluation in small bowel adenocarcinoma: analysis of the surveillance, epidemiology, and end results database. *Cancer* 2010;116:5374-5382. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20715162>.
59. Tran TB, Qadan M, Dua MM, et al. Prognostic relevance of lymph node ratio and total lymph node count for small bowel adenocarcinoma. *Surgery* 2015;158:486-493. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26013988>.
60. Wilhelm A, Muller SA, Steffen T, et al. Patients with adenocarcinoma of the small intestine with 9 or more regional lymph nodes retrieved have a higher rate of positive lymph nodes and improved survival. *J Gastrointest Surg* 2016;20:401-410. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26487334>.
61. Wu S, Chen JN, Zhang QW, et al. A new metastatic lymph node classification-based survival predicting model in patients with small bowel adenocarcinoma: A derivation and validation study. *EBioMedicine* 2018;32:134-141. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29908920>.
62. Zhang S, Yuan W, Zhang J, et al. Clinicopathological features, surgical treatments, and survival outcomes of patients with small bowel adenocarcinoma. *Medicine (Baltimore)* 2017;96:e7713. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28767610>.
63. Hashimoto D, Arima K, Chikamoto A, et al. Limited resection of the duodenum for nonampullary duodenal tumors, with review of the literature. *Am Surg* 2016;82:1126-1132. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28206943>.
64. Onkendi EO, Boostrom SY, Sarr MG, et al. 15-year experience with surgical treatment of duodenal carcinoma: a comparison of periampullary and extra-ampullary duodenal carcinomas. *J Gastrointest Surg* 2012;16:682-691. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22350721>.
65. Cloyd JM, Norton JA, Visser BC, Poultides GA. Does the extent of resection impact survival for duodenal adenocarcinoma? Analysis of 1,611 cases. *Ann Surg Oncol* 2015;22:573-580. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25160736>.



## NCCN Guidelines Version 2.2021 Small Bowel Adenocarcinoma

66. Raghav K, Overman MJ. Small bowel adenocarcinomas--existing evidence and evolving paradigms. *Nat Rev Clin Oncol* 2013;10:534-544. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23897080>.
67. U.S. National Library of Medicine. Phase III Trial Investigating the Potential Benefit of Adjuvant Chemotherapy for Small Bowel Adenocarcinoma (BALLAD). *ClinicalTrials.gov*: 2015. Available at: <https://clinicaltrials.gov/ct2/show/NCT02502370>. Accessed January 9, 2019.
68. Evans TRJ, Aparicio T, Malicot KL, et al. GLOBAL BALLAD: An International Rare Cancers Initiative trial to evaluate the potential benefit of adjuvant chemotherapy for small bowel adenocarcinoma (IRCI 002) [abstract]. *J Clin Oncol* 2016;34:TPS4154-TPS4154. Available at: [http://ascopubs.org/doi/abs/10.1200/JCO.2016.34.15\\_suppl.TPS4154](http://ascopubs.org/doi/abs/10.1200/JCO.2016.34.15_suppl.TPS4154).
69. Ecker BL, McMillan MT, Datta J, et al. Efficacy of adjuvant chemotherapy for small bowel adenocarcinoma: A propensity score-matched analysis. *Cancer* 2016;122:693-701. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26717303>.
70. Swartz MJ, Hughes MA, Frassica DA, et al. Adjuvant concurrent chemoradiation for node-positive adenocarcinoma of the duodenum. *Arch Surg* 2007;142:285-288. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17372054>.
71. Kelsey CR, Nelson JW, Willett CG, et al. Duodenal adenocarcinoma: patterns of failure after resection and the role of chemoradiotherapy. *Int J Radiat Oncol Biol Phys* 2007;69:1436-1441. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17689032>.
72. Aydin D, Sendur MA, Kefeli U, et al. Evaluation of prognostic factors and adjuvant chemotherapy in patients with small bowel adenocarcinoma who underwent curative resection. *Clin Colorectal Cancer* 2017;16:220-227. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27670893>.
73. Ye X, Zhang G, Chen H, Li Y. Meta-analysis of postoperative adjuvant therapy for small bowel adenocarcinoma. *PLoS One* 2018;13:e0200204. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30096150>.
74. Kim K, Chie EK, Jang JY, et al. Role of adjuvant chemoradiotherapy for duodenal cancer: a single center experience. *Am J Clin Oncol* 2012;35:533-536. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21659832>.
75. Overman MJ, Kopetz S, Lin E, et al. Is there a role for adjuvant therapy in resected adenocarcinoma of the small intestine. *Acta Oncol* 2010;49:474-479. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20397775>.
76. Ecker BL, McMillan MT, Datta J, et al. Adjuvant chemotherapy versus chemoradiotherapy in the management of patients with surgically resected duodenal adenocarcinoma: A propensity score-matched analysis of a nationwide clinical oncology database. *Cancer* 2017;123:967-976. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28263387>.
77. Markowitz SD, Bertagnolli MM. Molecular origins of cancer: Molecular basis of colorectal cancer. *N Engl J Med* 2009;361:2449-2460. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20018966>.
78. Kim GP, Colangelo LH, Wieand HS, et al. Prognostic and predictive roles of high-degree microsatellite instability in colon cancer: a National Cancer Institute-National Surgical Adjuvant Breast and Bowel Project Collaborative Study. *J Clin Oncol* 2007;25:767-772. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17228023>.
79. Klingbiel D, Saridaki Z, Roth AD, et al. Prognosis of stage II and III colon cancer treated with adjuvant 5-fluorouracil or FOLFIRI in relation to microsatellite status: results of the PETACC-3 trial. *Ann Oncol* 2015;26:126-132. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25361982>.
80. Ribic CM, Sargent DJ, Moore MJ, et al. Tumor microsatellite-instability status as a predictor of benefit from fluorouracil-based adjuvant chemotherapy for colon cancer. *N Engl J Med* 2003;349:247-257. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12867608>.
81. Sargent DJ, Marsoni S, Monges G, et al. Defective mismatch repair as a predictive marker for lack of efficacy of fluorouracil-based adjuvant



## NCCN Guidelines Version 2.2021 Small Bowel Adenocarcinoma

therapy in colon cancer. *J Clin Oncol* 2010;28:3219-3226. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20498393>.

82. Kim JE, Hong YS, Kim HJ, et al. Defective mismatch repair status was not associated with DFS and OS in stage II colon cancer treated with adjuvant chemotherapy. *Ann Surg Oncol* 2015;22 Suppl 3:S630-637. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26271397>.

83. Hutchins G, Southward K, Handley K, et al. Value of mismatch repair, KRAS, and BRAF mutations in predicting recurrence and benefits from chemotherapy in colorectal cancer. *J Clin Oncol* 2011;29:1261-1270. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21383284>.

84. Bertagnolli MM, Redston M, Compton CC, et al. Microsatellite instability and loss of heterozygosity at chromosomal location 18q: prospective evaluation of biomarkers for stages II and III colon cancer--a study of CALGB 9581 and 89803. *J Clin Oncol* 2011;29:3153-3162. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21747089>.

85. Compton CC, Fielding LP, Burgart LJ, et al. Prognostic factors in colorectal cancer. College of American Pathologists Consensus Statement 1999. *Arch Pathol Lab Med* 2000;124:979-994. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10888773>.

86. Fujita S, Shimoda T, Yoshimura K, et al. Prospective evaluation of prognostic factors in patients with colorectal cancer undergoing curative resection. *J Surg Oncol* 2003;84:127-131. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/14598355>.

87. Liebig C, Ayala G, Wilks J, et al. Perineural invasion is an independent predictor of outcome in colorectal cancer. *J Clin Oncol* 2009;27:5131-5137. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19738119>.

88. Grothey A, Sobrero AF, Shields AF, et al. Duration of adjuvant chemotherapy for stage III colon cancer. *N Engl J Med* 2018;378:1177-1188. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29590544>.

89. Onkendi EO, Boostrom SY, Sarr MG, et al. Neoadjuvant treatment of duodenal adenocarcinoma: a rescue strategy. *J Gastrointest Surg*

2012;16:320-324. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21956430>.

90. Yeung RS, Weese JL, Hoffman JP, et al. Neoadjuvant chemoradiation in pancreatic and duodenal carcinoma. A Phase II Study. *Cancer* 1993;72:2124-2133. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8374871>.

91. Coia L, Hoffman J, Scher R, et al. Preoperative chemoradiation for adenocarcinoma of the pancreas and duodenum. *Int J Radiat Oncol Biol Phys* 1994;30:161-167. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8083109>.

92. Minardi AJ, Jr., Zibari GB, Aultman DF, et al. Small-bowel tumors. *J Am Coll Surg* 1998;186:664-668. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9632155>.

93. Ciresi DL, Scholten DJ. The continuing clinical dilemma of primary tumors of the small intestine. *Am Surg* 1995;61:698-702; discussion 702-693. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/7618809>.

94. Ojha A, Zacherl J, Scheuba C, et al. Primary small bowel malignancies: single-center results of three decades. *J Clin Gastroenterol* 2000;30:289-293. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10777190>.

95. Oh SY, Edwards A, Mandelson M, et al. Survival and clinical outcome after endoscopic duodenal stent placement for malignant gastric outlet obstruction: comparison of pancreatic cancer and nonpancreatic cancer. *Gastrointest Endosc* 2015;82:460-468.e462. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25851162>.

96. van den Berg MW, Haijink S, Fockens P, et al. First data on the Evolution duodenal stent for palliation of malignant gastric outlet obstruction (DUOLUTION study): a prospective multicenter study. *Endoscopy* 2013;45:174-181. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23348890>.



# NCCN Guidelines Version 2.2021

## Small Bowel Adenocarcinoma

97. Upchurch E, Ragusa M, Cirocchi R. Stent placement versus surgical palliation for adults with malignant gastric outlet obstruction. *Cochrane Database Syst Rev* 2018;5:CD012506. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29845610>.
98. Cancer Stat Facts: Small Intestine Cancer. 2019. Available at: <https://seer.cancer.gov/statfacts/html/smint.html>. Accessed
99. Adam R, Chiche L, Aloia T, et al. Hepatic resection for noncolorectal nonendocrine liver metastases: analysis of 1,452 patients and development of a prognostic model. *Ann Surg* 2006;244:524-535. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16998361>.
100. Rompteaux P, Gagniere J, Gornet JM, et al. Resection of small bowel adenocarcinoma metastases: Results of the ARCAD-NADEGE cohort study. *Eur J Surg Oncol* 2019;45:331-335. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30501999>.
101. Rovers KP, de Bree E, Yonemura Y, de Hingh IH. Treatment of peritoneal metastases from small bowel adenocarcinoma. *Int J Hyperthermia* 2017;33:571-578. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27919181>.
102. Legue LM, Simkens GA, Creemers GM, et al. Synchronous peritoneal metastases of small bowel adenocarcinoma: Insights into an underexposed clinical phenomenon. *Eur J Cancer* 2017;87:84-91. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29132061>.
103. Elias D, Glehen O, Pocard M, et al. A comparative study of complete cytoreductive surgery plus intraperitoneal chemotherapy to treat peritoneal dissemination from colon, rectum, small bowel, and nonpseudomyxoma appendix. *Ann Surg* 2010;251:896-901. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20395843>.
104. Liu Y, Ishibashi H, Takeshita K, et al. Cytoreductive surgery and hyperthermic intraperitoneal chemotherapy for peritoneal dissemination from small bowel malignancy: Results from a single specialized center. *Ann Surg Oncol* 2016;23:1625-1631. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26717938>.
105. Liu Y, Yonemura Y, Levine EA, et al. Cytoreductive Surgery Plus Hyperthermic Intraperitoneal Chemotherapy for Peritoneal Metastases From a Small Bowel Adenocarcinoma: Multi-Institutional Experience. *Ann Surg Oncol* 2018;25:1184-1192. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29484565>.
106. Saxena A, Valle SJ, Liauw W, Morris DL. Recurrence and Survival Outcomes After Cytoreductive Surgery and Hyperthermic Intraperitoneal Chemotherapy for Small Bowel Adenocarcinoma. *Anticancer Res* 2017;37:5737-5742. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28982894>.
107. van Oudheusden TR, Lemmens VE, Braam HJ, et al. Peritoneal metastases from small bowel cancer: Results of cytoreductive surgery and hyperthermic intraperitoneal chemotherapy in The Netherlands. *Surgery* 2015;157:1023-1027. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25818658>.
108. Quenet F, Elias D, Roca L, et al. A UNICANCER phase III trial of hyperthermic intra-peritoneal chemotherapy (HIPEC) for colorectal peritoneal carcinomatosis (PC): PRODIGE 7 [abstract]. *J Clin Oncol* 2018;36:suppl; abstr LBA3503. Available at: <https://meetinglibrary.asco.org/record/158740/abstract>.
109. Czaykowski P, Hui D. Chemotherapy in small bowel adenocarcinoma: 10-year experience of the British Columbia Cancer Agency. *Clin Oncol (R Coll Radiol)* 2007;19:143-149. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17355111>.
110. Jigyasu D, Bedikian AY, Stroehlein JR. Chemotherapy for primary adenocarcinoma of the small bowel. *Cancer* 1984;53:23-25. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/6690001>.
111. Zhang L, Wang LY, Deng YM, et al. Efficacy of the FOLFOX/CAPOX regimen for advanced small bowel adenocarcinoma: a three-center study from China. *J BUON* 2011;16:689-696. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22331723>.



## NCCN Guidelines Version 2.2021 Small Bowel Adenocarcinoma

112. Aydin D, Sendur MA, Kefeli U, et al. Evaluation of prognostic factors and treatment in advanced small bowel adenocarcinoma: report of a multi-institutional experience of Anatolian Society of Medical Oncology (ASMO). *J BUON* 2016;21:1242-1249. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27837629>.
113. Pedersen K, Smyrk TC, Harrington S, McWilliams RR. Programmed death-ligand 1 (PD-L1) expression in small bowel adenocarcinomas (SBA) [abstract]. *J Clin Oncol* 2015;33:3619-3619. Available at: [http://ascopubs.org/doi/abs/10.1200/jco.2015.33.15\\_suppl.3619](http://ascopubs.org/doi/abs/10.1200/jco.2015.33.15_suppl.3619).
114. Thota R, Gonzalez RS, Berlin J, et al. Could the PD-1 pathway be a potential target for treating small intestinal adenocarcinoma? *Am J Clin Pathol* 2017;148:208-214. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28821192>.
115. Laforest A, Aparicio T, Zaan A, et al. ERBB2 gene as a potential therapeutic target in small bowel adenocarcinoma. *Eur J Cancer* 2014;50:1740-1746. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24797764>.
116. Takayoshi K, Kusaba H, Uenomachi M, et al. Suggestion of added value by bevacizumab to chemotherapy in patients with unresectable or recurrent small bowel cancer. *Cancer Chemother Pharmacol* 2017;80:333-342. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28653251>.
117. Gulhati P, Raghav K, Shroff R, et al. Phase II study of panitumumab in RAS wild-type metastatic adenocarcinoma of small bowel or ampulla of Vater. *Oncologist* 2018;23:277-e226. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29259073>.
118. Overman MJ, Varadhachary GR, Kopetz S, et al. Phase II study of capecitabine and oxaliplatin for advanced adenocarcinoma of the small bowel and ampulla of Vater. *J Clin Oncol* 2009;27:2598-2603. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19164203>.
119. Xiang XJ, Liu YW, Zhang L, et al. A phase II study of modified FOLFOX as first-line chemotherapy in advanced small bowel adenocarcinoma. *Anticancer Drugs* 2012;23:561-566. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22481063>.
120. Horimatsu T, Nakayama N, Moriwaki T, et al. A phase II study of 5-fluorouracil/L-leucovorin/oxaliplatin (mFOLFOX6) in Japanese patients with metastatic or unresectable small bowel adenocarcinoma. *Int J Clin Oncol* 2017;22:905-912. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28536826>.
121. Gibson MK, Holcroft CA, Kvols LK, Haller D. Phase II study of 5-fluorouracil, doxorubicin, and mitomycin C for metastatic small bowel adenocarcinoma. *Oncologist* 2005;10:132-137. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15709215>.
122. Tsushima T, Taguri M, Honma Y, et al. Multicenter retrospective study of 132 patients with unresectable small bowel adenocarcinoma treated with chemotherapy. *Oncologist* 2012;17:1163-1170. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22622149>.
123. Zaan A, Costes L, Gauthier M, et al. Chemotherapy of advanced small-bowel adenocarcinoma: a multicenter AGEO study. *Ann Oncol* 2010;21:1786-1793. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20223786>.
124. McWilliams RR, Foster NR, Mahoney MR, et al. North Central Cancer Treatment Group N0543 (Alliance): A phase 2 trial of pharmacogenetic-based dosing of irinotecan, oxaliplatin, and capecitabine as first-line therapy for patients with advanced small bowel adenocarcinoma. *Cancer* 2017;123:3494-3501. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28493308>.
125. Gulhati P, Raghav K, Shroff RT, et al. Bevacizumab combined with capecitabine and oxaliplatin in patients with advanced adenocarcinoma of the small bowel or ampulla of vater: A single-center, open-label, phase 2 study. *Cancer* 2017;123:1011-1017. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27859010>.
126. Aydin D, Sendur MA, Kefeli U, et al. Evaluation of bevacizumab in advanced small bowel adenocarcinoma. *Clin Colorectal Cancer*



## NCCN Guidelines Version 2.2021 Small Bowel Adenocarcinoma

2017;16:78-83. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27247089>.

127. Le DT, Uram JN, Wang H, et al. PD-1 blockade in tumors with mismatch-repair deficiency. *N Engl J Med* 2015;372:2509-2520. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26028255>.

128. U.S. Food and Drug Administration. FDA grants accelerated approval to pembrolizumab for first tissue/site agnostic indication. 2017. Available at: <https://www.fda.gov/Drugs/InformationOnDrugs/ApprovedDrugs/ucm560040.htm>. Accessed January 8, 2019.

129. Pedersen K, Foster N, Overman M, et al. ZEBRA: an ACCRU/IRCI multicenter phase 2 study of pembrolizumab in patients with advanced small bowel adenocarcinoma (SBA) [abstract]. *World Congress on Gastrointestinal Cancer 2019*; Abstract # O-007. Available at: <https://www.oncnet.com/meeting-materials/world-congress-gastrointestinal-cancer/3823>.

130. Overman MJ, McDermott R, Leach JL, et al. Nivolumab in patients with metastatic DNA mismatch repair-deficient or microsatellite instability-high colorectal cancer (CheckMate 142): an open-label, multicentre, phase 2 study. *Lancet Oncol* 2017;18:1182-1191. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28734759>.

131. Overman MJ, Lonardi S, Wong KYM, et al. Durable clinical benefit with nivolumab plus ipilimumab in DNA mismatch repair-deficient/microsatellite instability-high metastatic colorectal cancer. *J Clin Oncol* 2018;36:773-779. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29355075>.

132. Overman MJ, Adam L, Raghav K, et al. Phase II study of nab-paclitaxel in refractory small bowel adenocarcinoma and CpG island methylator phenotype (CIMP)-high colorectal cancer. *Ann Oncol* 2018;29:139-144. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29069279>.

133. Aldrich JD, Raghav KPS, Varadhachary GR, et al. Retrospective analysis of taxane-based therapy in small bowel adenocarcinoma. *Oncologist* 2018. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30598498>.

134. Zaanan A, Gauthier M, Malka D, et al. Second-line chemotherapy with fluorouracil, leucovorin, and irinotecan (FOLFIRI regimen) in patients with advanced small bowel adenocarcinoma after failure of first-line platinum-based chemotherapy: a multicenter AGEO study. *Cancer* 2011;117:1422-1428. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21425142>.

135. Drilon A, Laetsch TW, Kummar S, et al. Efficacy of Larotrectinib in TRK Fusion-Positive Cancers in Adults and Children. *N Engl J Med* 2018;378:731-739. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29466156>.

136. U.S. Food and Drug Administration. FDA approves larotrectinib for solid tumors with NTRK gene fusions. 2018. Available at: <https://www.fda.gov/Drugs/InformationOnDrugs/ApprovedDrugs/ucm626720.htm>. Accessed November 30, 2018.

137. Kim M, Jang HJ. The role of small bowel endoscopy in small bowel Crohn's disease: when and how? *Intest Res* 2016;14:211-217. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27433142>.

138. Simon M, Cosnes J, Gornet JM, et al. Endoscopic detection of small bowel dysplasia and adenocarcinoma in Crohn's disease: A prospective cohort-study in high-risk patients. *J Crohns Colitis* 2017;11:47-52. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27405958>.

139. Goverde A, Korsche SE, Wagner A, et al. Small-bowel surveillance in patients with Peutz-Jeghers syndrome: Comparing magnetic resonance enteroclysis and double balloon enteroscopy. *J Clin Gastroenterol* 2017;51:e27-e33. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27404294>.

140. Haanstra JF, Al-Toma A, Dekker E, et al. Prevalence of small-bowel neoplasia in Lynch syndrome assessed by video capsule endoscopy. *Gut*





# NCCN Guidelines Version 2.2021

## Small Bowel Adenocarcinoma

2015;64:1578-1583. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/25209657>.

141. Katsinelos P, Kountouras J, Chatzimavroudis G, et al. Wireless capsule endoscopy in detecting small-intestinal polyps in familial adenomatous polyposis. *World J Gastroenterol* 2009;15:6075-6079.

Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20027680>.

142. Koornstra JJ. Small bowel endoscopy in familial adenomatous polyposis and Lynch syndrome. *Best Pract Res Clin Gastroenterol* 2012;26:359-368. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/22704577>.

143. Postgate A, Hyer W, Phillips R, et al. Feasibility of video capsule endoscopy in the management of children with Peutz-Jeghers syndrome: a blinded comparison with barium enterography for the detection of small bowel polyps. *J Pediatr Gastroenterol Nutr* 2009;49:417-423. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/19543117>.

144. Serrano M, Mao-de-Ferro S, Pinho R, et al. Double-balloon enteroscopy in the management of patients with Peutz-Jeghers syndrome: a retrospective cohort multicenter study. *Rev Esp Enferm Dig* 2013;105:594-599. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24641456>.

145. van Lier MG, Wagner A, Mathus-Vliegen EM, et al. High cancer risk in Peutz-Jeghers syndrome: a systematic review and surveillance recommendations. *Am J Gastroenterol* 2010;105:1258-1264; author reply 1265. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20051941>.

146. Drini M, Speer A, Dow C, et al. Management of duodenal adenomatosis in FAP: single centre experience. *Fam Cancer* 2012;11:167-173. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/22131053>.

147. Hewitt M, Greenfield S, Stovall E, eds. *From Cancer Patient to Cancer Survivor: Lost in Transition*. Committee on Cancer Survivorship: Improving Care and Quality of Life, Institute of Medicine and National

Research Council: National Academy of Sciences; 2006. Available at:

<http://www.nap.edu/catalog/11468.html>.

148. Jansen L, Herrmann A, Stegmaier C, et al. Health-related quality of life during the 10 years after diagnosis of colorectal cancer: a population-based study. *J Clin Oncol* 2011;29:3263-3269. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/21768465>.

149. Lynch BM, Steginga SK, Hawkes AL, et al. Describing and predicting psychological distress after colorectal cancer. *Cancer* 2008;112:1363-1370. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18318044>.

150. Mols F, Beijers T, Lemmens V, et al. Chemotherapy-induced neuropathy and its association with quality of life among 2- to 11-year colorectal cancer survivors: results from the population-based PROFILES registry. *J Clin Oncol* 2013;31:2699-2707. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23775951>.

151. Thong MS, Mols F, Wang XS, et al. Quantifying fatigue in (long-term) colorectal cancer survivors: a study from the population-based patient reported outcomes following initial treatment and long term evaluation of survivorship registry. *Eur J Cancer* 2013;49:1957-1966. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23453750>.

152. Wright P, Downing A, Morris EJ, et al. Identifying social distress: a cross-sectional survey of social outcomes 12 to 36 months after colorectal cancer diagnosis. *J Clin Oncol* 2015;33:3423-3430. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/26282636>.

153. Denlinger CS, Barsevick AM. The challenges of colorectal cancer survivorship. *J Natl Compr Canc Netw* 2009;7:883-893; quiz 894.

Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19755048>.